

An extended frame for cosmology by integrable Weyl geometry

Erhard Scholz,¹

Wuppertal University

Department C, Div. of Mathematics

42097 Wuppertal, Germany

Abstract

A conservative extension of general relativity by integrable Weyl geometry is formulated, and a new class of cosmological models (*Weyl universes*) is introduced and studied. A short discussion of how these new models behave in the light of observational evidence follows. It turns out that the subclass of Weyl universes with positive ex-ante curvature of spatial sections (*Einstein-Weyl universes*) forms an empirically contentful conceptual frame for cosmology. Recent high precision data (supernovae Ia, quasar frequencies, microwave background) are well represented in this type of Weyl geometry and several of the anomalies of the standard approach are avoided or become meaningless.

Introduction

Present standard cosmology relies on the physical geometry of Friedman-Lemaître models and draws upon the hypothesis of expanding spatial sections in relativistic space-time. For several decades, this approach was a satisfying conceptual frame for astrophysical and astronomical research. Even though many continue to consider it as a reliable guide for the evaluation of empirical observations, there are strong indications of growing anomalies for this interpretational scheme, most prominent among them the appearance of “non-baryonic dark matter”, the surprising “time-dependence of vacuum (dark) energy”) and the “scale problem of vacuum energy”. Other problems, like the isotropy/anisotropy riddle of the cosmic microwave background, are covered up by surprising towers of ad-hoc constructions or are neglected, like the negative results on a systematic correlation between redshift and metallicity of galaxies and quasars. It is not the goal of this paper to discuss these questions in detail. We rather want to show that already a small and innocent looking modification in the conceptual framework of Robertson-Walker manifolds which underly the geometry of the Friedman-Lemaître models (F-L), allows to conceive of new physical geometries which are consistent with general relativity (GRT).² They can be well adapted to observational data and give an account of several of the crucial tests for cosmological models,

¹scholz@math.uni-wuppertal.de

²For a general survey of non-Riemannian geometries as possible alternatives for cosmological geometry see (Puetzfeld 2004).

at least as convincingly as in the F-L approach (redshift-distance relationship, SN_{Ia} magnitudes). In some respects, they are even more convincing (redshift dependence of quasar frequencies and metallicity), in others new possibilities are open for research (microwave background). Above all, no dynamical “dark energy” anomaly appears in these models. The right hand side of the Einstein equation looks much more realistic than in the expanding world approach, although also here questions with respect to the state equation of the cosmic medium remain.

Since the 1920s, when cosmological redshift started to be observed and became the first empirical clue to the construction of cosmological geometries backed on general relativity, some of the leading protagonists, including E. Hubble himself, could never be convinced that the space-kinematical interpretation of cosmological redshift was more than a provisional move to characterize the energy reduction of photons transmitted over long cosmic distances, while in the future a “more physical explanation” (H. Weyl in 1930) might be found. Different proposals for “tired light” approaches, as they have been called by their critics at first, have been tried, among them the ideas of a “gravitational drag” of photons (F. Zwicky, MacMillan e.a.) or of a quantum mechanical photon-photon interaction (E. Finlay-Freundlich, M. Born, J.-P. Vigi er e.a.).³ Although these investigations have not acquired a physically reliable, in some sense “verified” status, they contributed to a first exploration of possible physical causes of such a hypothetical mechanism. With the presently maturizing crisis in standard cosmology, it seems more than justified to understand better, how such hypothetical explanations can be conceptually integrated into general relativistic cosmology and to explore their mathematical relationship with the standard approach. The result of such investigations may contribute to a change of the symbolic “a-priori” (in a technical sense) of the evaluation of empirical data in observational cosmology. If not, they should at least add to our understanding of the expanding space cosmologies from “outside”, i.e., without assuming an expansion as an ex-ante property of the models.

In this article, we analyze cosmological redshift in Robertson-Walker models in terms of (scale-integrable) Weyl geometry. This allows to refrain from any ex-ante decision as to the physical explanation of cosmological redshift, either as space-kinematical (“Doppler”) or as tired light. It even allows us, to a certain degree, to translate mathematically between the two. We will see that there are not only those Robertson-Walker models which have a convincing physical interpretation under the assumption of a “real” space-expansion (the classical Friedman-Lema tre models), but also others which allow a convincing physical interpretation under the assumption of a tired light hypothesis. There is a particularly simple class of physically mean-

³For Finlay-Freundlich see (Hentschel 1998, 634ff.) and (Hentschel 1994, 187ff.) or the less critical report (including remarks on Born) (Assis/Neves 1995).

ingful Robertson-Walker models of the new type, which will be called *Weyl universes*, because they rely essentially on simple aspects of Weyl geometry.⁴ It is possible to interpret the data of observational cosmology in both model classes and to see in which approach physico-geometrical reality appears more convincing. That cannot be done here in detail, although some indications on first results of empirical evaluations (“tests” of the models) will be given at the end of this paper.

1. Weyl geometry

H. Weyl’s generalization of Riemannian and semi-Riemannian geometry, first presented in (Weyl 1918), is still being discussed in the actual literature, although the original intentions of its inventor for its physical application have not been realized. For mathematicians, Weyl geometry became an interesting structure after fibre bundles had been invented and Weyl’s idea could be rephrased in this language (Folland 1970, Varadarajan 2003). In the physical literature it played an important part in the first wave of unified field theories and seemed to loose its importance with it.⁵ But it contained much broader possibilities than its direct semantical link to Weyl’s first (metrical) gauge theory of electromagnetism made visible. In his “mathematical analysis of the space problem” during the years 1921–23, Weyl gave strong arguments in this direction.⁶ The conceptual potential was taken up again, in a different sense, by J. Ehlers, F. Pirani and A. Schild in their constructive axiomatic approach to general relativity (Ehlers e.a. 1972). Thus the question arose, whether the additional structural element of Weyl geometry, the length connection φ , could play the role of some, perhaps still unknown, field. A negative answer was given by J. Audretsch, F. Gähler and N. Straumann, who showed that a Weyl geometry can only be considered as compatible with relativistic quantum mechanics, if the length connection is integrable (Audretsch e.a. 1984). Such a restriction had already been proposed by P.A.M. Dirac in his (Dirac 1973). Field theoretically, the extension of GRT by Weyl geometry appeared thus to be condemned to triviality.

This is not the end of the story, however; see, e.g., (Tiwari 1989, Tiwari 2003). Here we reconsider Weyl geometry from a genuinely geometrical perspective which is, in its metrical aspects, very close to Weyl’s original view. In this section the aspects of Weyl geometry which are most important for us will be resumed.⁷ In the next one we see how it can be used for a weak,

⁴They were *not* already introduced by H. Weyl, although he could easily have done so, had he broken with the dogma of the electromagnetic potential of his gauge geometry early enough.

⁵See (Goenner 2004).

⁶See (Scholz 2004b).

⁷Although there is considerable overlap with section 3 of (Dirac 1973), readers acquainted with Dirac’s formalism should be aware that Dirac switched sign in the transformation of the length connection, in our notation $\tilde{\varphi} = \varphi + d \ln \lambda$ rather than $\tilde{\varphi} = \varphi - d \ln \lambda$

i.e. conservative, extension of general relativity, even if the restriction of Audretsch, Gähler and Straumann is taken into account. In the third and the following sections, a method is developed which allows to use integrable Weyl geometry for a generalized view on Robertson-Walker manifolds and for cosmological model building.

Weyl proposed to found the metrical concepts of differential geometry on a “purely infinitesimal” point of view more strictly than in Riemannian geometry. In particular, he was not satisfied with the possibility, inbuilt into classical differential geometry, to compare lengths of vectors at different points of a manifold directly. He considered it as more natural to compare lengths of vectors directly, only if they are attached to the same point, $\xi, \eta \in T_p M$ for some $p \in M$. That can easily be realized by a specification of a conformal semi-Riemannian structure on M . For the comparison of lengths of vectors at different points $p, q \in M$, $\xi \in T_p M$, $\eta \in T_q M$, Weyl introduced a *length connection*, given by a differential 1-form φ on M . This form was used to define a kind of “length transfer” by integration along paths, comparable to the parallel transfer of vectors by an affine connection. Of course, other geometrical or physical quantities related to length or time measurement (“gauge quantities” here introduced as Weyl fields in definition 5) have to be compared in a similar manner. The change of the representative of the conformal metric leads to a change of the differential 1-form of the length connection (Weyl 1923, 122ff.). The result of Weyl’s rearrangement of basic concepts of differential geometry can be stated in the following terms:⁸

Definition 1 *a) A gauge of a differentiable manifold M is given by a pair (g, φ) consisting of a semi-Riemannian metric g on M of any signature (p, q) (here usually of Lorentzian signature $(p, q) = (1, 3)$) and a real-valued differential 1-form φ on M . The 1-form φ will be called the **length connection** (or **scale connection**) of the gauge and g its **Riemannian component**.
b) Two gauges (g, φ) and $(\tilde{g}, \tilde{\varphi})$ are considered as **gauge equivalent**, iff there is a strictly positive real valued function λ on M , such that*

$$\tilde{g} = \lambda^2 g, \quad (1)$$

$$\text{and } \tilde{\varphi} = \varphi - d \ln \lambda = \varphi - \frac{d\lambda}{\lambda}. \quad (2)$$

*A transformation between gauge equivalent gauges will be called a change of gauge, with **gauge factor** (or a rescaling by the **scale transformation**) λ and **gauge transformation** (2).*

*c) An equivalence class of gauges, denoted by $[g, \varphi]$, is called a **Weylian** (Dirac 1973, eqs. (1.2), (1.3)). This change destroys the behaviour as a proper gauge transformation in the sense of differential geometry (footnote 9). In particular, Weyl’s gauge invariant comparison of length ratios (our lemma 1) gets lost.*

⁸The formulations of this passage owes much to proposals by M. Kreck. Compare also (Folland 1970, Varadarajan 2003).

metric on M and a manifold with Weylian metric $(M, [g, \varphi])$ will be called a Weylian manifold.

*d) A gauge will be called **semi-Riemannian**, if it is of the form $(g, 0)$, i.e., its length connection is zero. A Weylian manifold will be called **semi-Riemannian**, if it has a semi-Riemannian gauge.*

Remark 1 If one considers the trivial principal bundle $\mathcal{S}M \longrightarrow M$ with group $(\mathbb{R}^+, \cdot)$ and total space $M \times \mathbb{R}^+$, a connection is the same as a real-valued 1-form. Here the gauge group $\Gamma(\mathcal{S}M)$ is the set of (differentiable) maps $M \longrightarrow \mathbb{R}^+$. $\mathcal{S}M$ will be called the *scale bundle*. The group $\mathbb{R}^+$ operates on the connections as given in (2). In order to define a Weylian metric, one has to specify a pair consisting of a connection in $\mathcal{S}M$ and a metric g on M , and to introduce an equivalence relation as in b). Equivalently, if one starts from a conformal structure $[g]$ on M , represented by g , a Weylian metric is specified by the selection of a connection in the scale bundle. Change of representatives of $[g]$ go along with changes of the trivialization of $\mathcal{S}M$ and the corresponding gauge transformation for the 1-form. This view turns out to be useful when conformal approaches to cosmology or field theory are related to the Weyl geometric point of view.

Here we do not need the bundle structure and we work with our definition 1, which is introduced very much in the spirit of Weyl's original presentation. It may be worthwhile to mention that the "gauge transformations" of our definition are *gauge transformations* of connections in $\mathcal{S}M$ in the sense of modern differential geometry.⁹ If other gauge structures enter the game, one ought to add the attribute "scaling", to make clear which (part of an perhaps extended) gauge group is referred to. In fact, the terminology "gauge" transformation is historically derived from this exemplar.

In the sense of remark 1, the curvature F of a length connection is well defined. If the connection is given by the differential form φ , the curvature is simple to calculate; it is nothing but the exterior differential

$$F = d\varphi. \quad (3)$$

F is obviously gauge independent and will be called the *length curvature* or *scale transfer curvature* of the Weylian manifold.

For the sake of notational simplicity, in the sequel the symbol g will often be used as a shorthand expression for the local matrix (g_{ij}) of g . Of course the expression of g could just as well be related to an orthonormal frame (ONF) description of the tangent bundle. That is advisable in contexts

⁹After local trivialization, the connection can be expressed by a Lie algebra valued 1-form A ; a change of gauge is expressed by the operation of a Lie group valued function γ . Because of commutativity the ensuing gauge transformation for the 1-form, $\tilde{A} = \gamma A \gamma^{-1} + \gamma d\gamma^{-1}$, reduces here to $\tilde{\varphi} = \varphi + \lambda d(\lambda^{-1}) = \varphi - d \ln \lambda$. Of course, the sign is important for the consistency of gauge geometrical considerations, cf. footnotes 7, 10.

where physical reference frames are necessary in the theoretical construction, as for the description of the general relativistic Dirac equation (Weyl 1929, Audretsch e.a. 1984) etc.. The geometry of Weylian manifolds will be called Weylian *gauge geometry* (in the original sense of the word), or in short *Weyl geometry*.

Remark 2 An *isomorphism* of Weylian manifolds $(M, [g, \varphi])$ and $(\tilde{M}, [\tilde{g}, \tilde{\varphi}])$ is given by a diffeomorphism $\beta : M \rightarrow \tilde{M}$, which transforms the Weylian metrics into each other,

$$\beta_* \tilde{g} = \lambda g, \quad \beta_* \tilde{\varphi} = \varphi - d \ln \lambda,$$

where λ is a gauge factor and β_* is an abbreviated notation for the linearization of β operating on differential forms (of any degree). Clearly λ depends on the choice of gauges in M and $\tilde{M}$. The defining data are $(\beta, [\lambda])$.

According to Weyl's argumentation of 1918, any representative g of the conformal metric may be used for a direct metrical comparison of vector or tensor data at one and the same point $p \in M$, or in its infinitesimal neighbourhood. The comparison of metrical data given at different points p and q ("of finite distance") can be achieved only indirectly (Weyl 1918, Weyl 1923, etc.). The length connection φ can be used to "transport length units" along a (a sufficiently differentiable) curve $c(t)$. More formally let us introduce:

Definition 2 *With respect to the gauge (g, φ) , a **length transfer or scale transfer** along a differentiable curve $c : [0, 1] \rightarrow M$ is defined as the real valued function $l : [0, 1] \rightarrow \mathbb{R}^+$ with*

$$l(t) := l(q) = e^{\int_0^t \varphi(c'(\tau)) d\tau}. \quad (4)$$

Remark 3 As l is not dependent on the parametrization, we often write $l(p) := l(c(t))$, if a curve c is specified and $p = c(t)$. A length transfer makes metrical quantities at $p = c(0)$ and $q = c(t)$ comparable, although its definition depends on the gauge and the path. That is shown by the next lemma. One could equivalently define the length transfer by a differential equation

$$l'(t) = l(t) \varphi(c'(t)), \quad l(0) = 1.$$

Weyl preferred a differential point of view and expressed this condition in terms of the change of the "infinitesimal length factor" dl

$$dl = l \sum_i \varphi_i dx^i.$$

Transformation under change of gauge is given by

$$\tilde{l} = e^{\int (\varphi - d \log \lambda)} = e^{-\log \lambda} e^{\int \varphi} = \lambda^{-1} l. \quad (5)$$

An immediate consequence is the following observation.

Lemma 1 For vectors $\xi \in T_p M$ and $\eta \in T_q M$ and l the length transfer with respect to a gauge (g, φ) , the modified quotient

$$l(c(t))^2 \frac{g_{c(t)}(\eta, \eta)}{g_p(\xi, \xi)}$$

is independent of the gauge. It may be used for a gauge invariant comparison of lengths of vectors at different points of a Weylian manifold.

For a **proof** one only needs to notice that the gauge transformations for the terms l^2 and g cancel.¹⁰

In general the comparison depends on the choice of the path. For the case of integrable Weyl manifolds (see below), to which we will restrict our view from the following section onward, it does not. In any case, the length transfer can be used as a factor to modify the measurement of vectors along a curve in a given metrical component g of a gauge (g, φ) ; i.e., one measures the length by $l^2 g$ rather than by g and arrives at a gauge invariant concept of comparison.

Remark 4 Weyl called this comparison of vectors at different points “calibration by transfer”. Its invariance under changes of gauge was behind his choice of terminology “gauge transformation” in the literal sense of the word. If a curve c with $c(0) = p$ and $c(1) = q$ is given and two equivalent gauges (g, φ) , $(\tilde{g}, \tilde{\varphi})$, with length transfer factors l and $\tilde{l}$ respectively (along c), are related by a gauge transformation with gauge factor λ , $\tilde{g} = \lambda^2 g$, the above equation (5) is written more precisely as

$$\tilde{l}(p, q) = \frac{\lambda(p)}{\lambda(q)} l(p, q).$$

Thus length transfer functions are changed under a gauge transformation by a gauge factor $\lambda(p)$ in the first variable and by $\lambda^{-1}(q)$ in the second one. In Weyl’s terminology which will be explained in a moment (definition 5), length transfer is given by a function $l(p, q)$ of *gauge weight* +1 in the first variable and −1 in the second.

Weyl gave strong arguments that his generalized metric results from an analysis of how one may understand metrical relationships from a “purely infinitesimal” point of view and that it is a better conceptual starting point than Riemannian geometry for the development of physical geometry in the sense of general relativity.

He considered it as a clue to his gauge geometry that a Weylian manifold has a uniquely determined compatible affine connection (*fundamental theorem* of Weyl geometry). *Compatibility* means that parallel transfer of vectors

¹⁰With Dirac’s convention for “gauge” transformations (cf. footnote 7) the gauge weight of length transfer changes sign and the lemma no longer holds.

preserves angles and that the length of parallel transported vectors changes with the length transfer along the curve (which is gauge invariant by Weyl's lemma 1).

Theorem 1 (Weyl) *A Weylian manifold $(M, [g, \varphi])$ has a uniquely determined affine connection Γ compatible with the Weylian metric. If a gauge (g, φ) is chosen, and ${}_g\Gamma$ is the Levi-Civita connection of g alone, the following relation holds:*

$$\Gamma_{jk}^i = {}_g\Gamma_{jk}^i + (\delta_j^i \varphi_k + \delta_k^i \varphi_j - g_{jk} \varphi^i)$$

(using Einstein sum convention for common coordinate indexes in the given quantities and the Kronecker symbol δ).

Clearly ${}_g\Gamma$ is gauge dependent, while Γ is not. For a **proof** see, e.g., (Weyl 1923, 124f.). The theorem is behind the following

Definition 3 *The uniquely determined compatible affine connection Γ of a Weylian manifold $(M, [g, \varphi])$ will be called the **Weyl-Levi-Civita connection**, or in short **W-L-C connection**, of M . By covariant derivation in M we refer to the derivation $D = D_\Gamma$ with respect to the W-L-C connection, if not otherwise specified.*

As the Riemann curvature tensor of type (1.3), $R = (R_{jkl}^i)$, can be geometrically defined from the affine connection only (without reference to the metric), the Riemann curvature of the W-L-C connection is independent of gauge. The same holds for the Ricci curvature as its contraction (which does not involve the Riemannian component g of the metric), whereas scalar curvature $\bar{R}$ as the contraction of $R_j^i = g^{ik} R_{kj}$, with $(R_{jk}) = Ric$ is obviously gauge dependent (we will see in a moment that $\bar{R}$ is a scalar function of “gauge weight” -2). Let us resume this observation as

Lemma 2 (Weyl) *The Riemann curvature of type (1,3), $R = (R_{jkl}^i)$, and the Ricci curvature of the Weyl-Levi-Civita connection are well defined tensor fields (i.e. gauge independent) on a Weylian manifold.*

It is therefore useful to agree upon the following terminology:

Definition 4 *In a Weylian manifold $(M, [g, \varphi])$ with Weyl-Levi-Civita connection Γ we define:*

- a) *The **Riemann** respectively **Ricci** curvature of the Weylian metric are the respective curvature tensors R and $Ric = (R_{ij})$ of Γ .*
- b) *The **scalar curvature** $\bar{R}$ is a family of scalar functions $\bar{R}_{(g, \varphi)}$ on M , indexed by the gauges (g, φ) , defined as $\bar{R}_{(g, \varphi)} := g^{ij} R_{ij}$ (using Einstein's sum convention).*
- c) *The **length curvature** or **scale transfer curvature** F is the (gauge independent) exterior differential $F = d\varphi$ for any gauge of M (see equ. (3)).*

The integration of the infinitesimal length transfer (4) is independent of the path, iff the curvature of the length connection vanishes. That gives an easy criterion for the possibility to find a semi-Riemannian gauge of a Weylian manifold.

Theorem 2 (Weyl) *A Weylian manifold $(M, [g, \varphi])$ is locally semi-Riemannian, iff its length curvature vanishes,*

$$F = d\varphi = 0.$$

Proof In this case, a semi-Riemannian gauge of M can be calculated from an arbitrary gauge (g, φ) by the path independent integrals starting from a reference point p (in any path connected component of M) and a path $c : [0, 1] \rightarrow M$ from p to q

$$\lambda(q) = e^{\int_0^1 \varphi(c'(\tau)) d\tau}. \quad (6)$$

As $d\varphi = 0$, the length transfer function is path independent in simply connected regions and can be used as a gauge factor. This leads to a $\tilde{g} = \lambda^2 g$ and $\tilde{\varphi}(q) = \varphi(q) - d \ln \lambda(q) = \varphi(q) - \varphi(q) = 0$ and thus to a semi-Riemannian gauge. Two semi-Riemannian gauges differ only by constants on each path-wise connected component of M .

Vanishing of the length curvature is an integrability condition and decides, whether a Weylian manifold is semi-Riemannian. In the case it is, we will call the Weyl geometry *integrable*. Although for integrable Weyl manifolds it seems at first glance that we can just as well go over to the view of semi-Riemannian geometry, we shall see that Weyl geometry allows to consider physically non-trivial metrical modifications of semi-Riemannian geometry *even in this special case*. Audretsch, Gähler and Straumann have shown that there are strong reasons to restrict attention to integrable Weyl manifolds in the physical context (Audretsch e.a. 1984). We will come back to their argument in the next section. Mathematically their argument builds on the following observation for Weyl manifolds:

Lemma 3 (Audretsch/Gähler/Straumann) *If a Weylian manifold $(M, [g, \varphi])$ with W-L-C connection Γ has a scalar Weyl field f with vanishing covariant derivative,*

$$D_\Gamma f = 0,$$

M is integrable.

Proof If the scalar Weyl field is given by f in some gauge (g, φ) , the covariant derivative is

$$D_\Gamma f = df - f \varphi,$$

where df denotes the ordinary differential. Vanishing of the covariant derivative implies

$$\varphi = \frac{df}{f} = d \log f .$$

Thus the length connection is a complete differential and integrable by the scale function $\lambda := f$.

If a tensor or a tensor field X is defined mathematically or empirically in any way by some relationship to metrical quantities, it may change under a change of gauge. Therefore we have to take into account the gauge behaviour of vectors, tensors, and their densities, if we want to do physics in the generalized geometrical framework (Weyl 1918, Weyl 1923). In order to give a well defined mathematical meaning to this concept, we introduce:

Definition 5 *a) Let (g, φ) , $(\tilde{g}, \tilde{\varphi})$ be gauges in a Weylian manifold $(M, [g, \varphi])$ and $X, \tilde{X}$ tensor fields (or tensor densities) on M . We then define triples to be **k-equivalent**, $(g, \varphi, X) \sim_k (\tilde{g}, \tilde{\varphi}, \tilde{X})$, iff the gauges (g, φ) , $(\tilde{g}, \tilde{\varphi})$ are gauge equivalent with gauge factor λ , i.e. $\tilde{g} = \lambda^2 g$, and*

$$\tilde{X} = \lambda^k X.$$

*b) A “gauge quantity” or a **Weyl field of gauge weight k** is a k -equivalence class of tripels $[g, \varphi, X]_k$.*

To simplify language, we often use representatives X of the third partner of the tripel and speak of tensors of weight k , if the context makes clear which gauge (g, φ) we refer to. Covariant differentiation of Weyl fields contain terms dependent on the gauge weight. We do not use it here and refer *in this respect* to (Dirac 1973, sec. 3).

Remark 5 For the sake of brevity, let us use double square brackets to indicate the gauge weight of geometrical or physical quantities, e.g. $[[g_{ij}]] = [[g]] = 2$, as the metric field $X = g$ has gauge weight 2 by definition; similarly $[[g^{ij}]] = -2$. We know already that the Ricci tensor of a Weylian manifold is gauge invariant, i.e., its gauge weight is $[[Ric]] = 0$. The same holds, of course, for all “classical” (gauge invariant) tensor (including vector and scalar) fields or densities. The scalar curvature $\bar{R}$ of a Weylian manifold is a scalar function of gauge weight $[[\bar{R}]] = -2$, as can be seen from the relationship $\bar{R} = g^{ij} R_{ij}$ ($Ric = R_{ij}$ in coordinate notation).

It is obvious how to form tensor products of Weyl fields of gauge weights k and l . Clearly the product is a Weyl field of weight $k + l$. Thus, although neither the Riemannian component g of a Weylian metric $[g, \varphi]$ nor its scalar curvature $\bar{R}$ are gauge invariant, their product is (gauge weight $2 - 2 = 0$). Therefore $\bar{R}g$ can be considered as a “classical”, i.e., gauge invariant, tensor field.

Time, the measure dimension of which will be denoted as $[t]$, has gauge weight $+1$, as have lengths $[l]$, whereas energy as its conjugate quantity (measure dimension $[E]$) is of gauge weight $[[E]] = -1$. Energy density is thus of gauge weight $[[E]][l^{-3}] = -1 + (-3) = -4$ and so on.¹¹

The concept of length transfer can now be generalized to scalar Weyl fields of arbitrary gauge weight. Weyl introduced a suggestive terminology (“calibration by transfer”) which we want to adopt in the following sense:

Definition 6 *If l is the length transfer function along a curve $c : [0, 1] \rightarrow M$ with respect to a gauge (g, φ) , a scalar Weyl field f of gauge weight k , defined along c , is said to propagate by **calibration transfer** along c , iff for any pair of values $0 \leq a \leq b \leq 1$ and $c(a) = p$, $c(b) = q$*

$$f(q) = \left(\frac{l(q)}{l(p)} \right)^k f(p). \quad (7)$$

An easy calculation shows that it is possible to gauge a Weylian manifold with nowhere vanishing scalar curvature, $\bar{R} \neq 0$ everywhere, such that the representative of scalar curvature becomes constant, e.g., $\bar{R} = \pm 1$. Starting from any gauge (g, φ) , this condition can easily be satisfied by applying the gauge factor $\lambda^2 = |\bar{R}|$. This gauge transformation gives

$$\tilde{g} = |\bar{R}|g = \lambda^2 g, \quad \tilde{\varphi} = \varphi - d \ln \sqrt{|\bar{R}|}$$

and leads to $\tilde{\bar{R}} = \lambda^{-2} \bar{R} = \frac{\bar{R}}{|\bar{R}|} = \pm 1$, as wanted.

Lemma 4 (Weyl) *For a Weylian manifold $(M, [g], \varphi)$ of nowhere vanishing scalar curvature $\bar{R}$, there is a gauge which normalizes scalar curvature to $\bar{R} = \text{const}$. Of course, such a gauge is unique up to a constant.*

Weyl assigned this particular gauge a particularly important role in his explorations of the use of Weyl geometry in cosmology, and we will follow and continue this track. We therefore introduce the terminology:

Definition 7 *In a Weylian manifold $(M, [g], \varphi)$ of nowhere vanishing scalar curvature, a gauge with constant scalar curvature will be called a **Weyl gauge** of M .*

Example 1 Consider a *Robertson-Walker manifold* $M = I \times_f S_\kappa$ with fibre S_κ over an open interval $I \subset \mathbb{R}$ with respect to a warp function $f : I \rightarrow \mathbb{R}^+$, where S_κ is a 3-dimensional Riemannian manifold with metric $d\sigma^2$ of

¹¹Weyl considered the gauge factor of the metric itself as the main reference for weights. His gauge weights are therefore half of the ones used here. The latter are closer to common dimensional considerations of physical quantities (see, e.g., (Manin 1981)).

constant sectional curvature $\kappa = \frac{k}{a^2}$ ($k = \pm 1$ or 0 and $a > 0$) (O'Neill 1983). Using the parameter τ for elements in I , the Lorentzian warp metric $\tilde{g}$ on M can be written as

$$\tilde{g}: ds^2 = d\tau^2 - (fd\sigma)^2.$$

If we introduce the Weylian metric by the gauge $(\tilde{g}, \tilde{\varphi})$ with $\tilde{\varphi} = 0$, we call $(M, [\tilde{g}, 0])$ a *Robertson-Walker-Weyl manifold*. We easily change the gauge to (g_1, φ_1) , such that one gets “static rest spaces”, i.e., constant metric along the spatial fibres

$$g_1 := f^{-2}\tilde{g} = \frac{d\tau^2}{f^2} - d\sigma^2, \quad \varphi_1 = -d\ln f^{-1} = \frac{df}{f}.$$

One should *not be deceived* to think that now the spatial sections might be gauged to constant curvature in Weyl geometry. That cannot be the case, in general, as sectional and scalar curvatures are of gauge weight -2 and we know the values of these curvature parameters for Robertson-Walker manifolds in the semi-Riemannian gauge. The scalar curvature $\tilde{R}$ and sectional curvatures $\tilde{\kappa}_T$ and $\tilde{\kappa}_S$ in plane directions containing the vector $\frac{\partial}{\partial\tau}$ (“cosmic timelike”) or in plane directions tangential to the spatial fibres respectively are in semi-Riemannian gauge $(\tilde{g}, 0)$ (see e.g. (O'Neill 1983, 345)):

$$\begin{aligned} \tilde{R} &= 6 \left(\left(\frac{f'}{f} \right)^2 + \frac{\kappa}{f^2} + \frac{f''}{f} \right), \\ \tilde{\kappa}_T &= \frac{f''}{f}, \quad \tilde{\kappa}_S = \left(\frac{f'}{f} \right)^2 + \frac{\kappa}{f^2}. \end{aligned}$$

We thus arrive at a *Weyl gauge* (g, φ) by setting

$$g := |\tilde{R}|g, \quad \varphi = -\frac{1}{2}d\ln |\tilde{R}|.$$

Now we have constant scalar curvature. To achieve constant sectional curvatures (in cosmic timelike *or* in spacelike direction) one has to use the respective values of sectional curvatures as scale gauge factors.

In the case of a Weylian manifold of vanishing length curvature and nowhere vanishing scalar curvature, we now have two distinguished gauges, *Riemann gauge* ($\varphi = 0$) and *Weyl gauge* ($\bar{R} \equiv \text{const}$). Even in the last edition of *Raum - Zeit - Materie*, two years after Weyl had given up his belief in the physical reality of his early unified field theory, he insisted upon the utility to study the interplay of both gauges. He guessed that it might be particularly useful for the understanding of a hypothetical “cosmological term” in the (generalized) Einstein equation and might contribute to an understanding of cosmological redshift (Weyl 1923, 300ff.).

2. A conservative extension of GRT

Until about 1926/27, Weyl tended to bind the usage of his gauge geometry in physics to an interpretation of the length connection φ as the electromagnetic potential. Let us call this interpretation the *e.m.p.-dogma* of gauge geometry. It was not the only possibility, even at the time. In fact, Weyl attempted to explore the possibilities opened up by gauge geometry for an understanding of cosmology, in particular redshift, at different occasions. These remarks stood in tension, and sometimes even in sharp conceptual dissonance, with the e.m.p.-dogma. By several reasons Weyl never explored the possibilities of his gauge geometry for cosmology in a “pure” form, i.e., *separated* from the e.m.p.-dogma.

Two scientific generations later, the dogma was broken. J. Ehlers, F. Pirani, and A. Schild had a different perspective, when they used Weyl geometry for a conceptual foundation of GRT (“constructive axiomatics” in the terminology of philosophers of science) (Ehlers e.a. 1972). They showed that under very convincing simple assumptions for the propagation of light and for the free fall of test particles, which are minimal conditions for the validity of special relativity in the infinitesimal approximation, a conformal Lorentzian structure is specified (null geodesics as descriptors for the light ray structure). The trajectories of freely falling test particles determine a projective class of affine connections (trajectories pass along the geodesics). If both sets of data are compatible (null geodesics of the light structure are a subclass of the affine geodesics), the structure of a Weylian manifold is obtained. Notwithstanding their foundational intentions, Ehlers e.a. left it open, whether Weyl’s length curvature F and the corresponding conserved current associated to the gauge symmetry $\mathbb{R}^+$ might perhaps relate the gravitational field to another universally conserved current and may thus “contain some physical truth” (Ehlers e.a. 1972, 83). About that time, high energy physicists had come to the conclusion that deep elastic scattering of electrons at nucleons was approximatively invariant under mass/energy scaling, and Weyl’s “length” gauge arose new interest among physicists. Now the scale invariance and its breaking by a scale connection was reconsidered (Deser 1970, Canuto e.a. 1977, Hehl e.a. 1989).

Audretsch, Gähler and Straumann damped too high expectations in this respect. They investigated Dirac and Klein-Gordon matter fields on a Weylian manifold M (assuming that the topological obstruction for existence of a spinor bundle over M vanishes). In both cases a mass function m on M appears in the equations, which acquires gauge weight -1 in the Weyl geometry setting. (In our terminology m is a scalar Weyl field of weight -1 .) They considered the series development of the solutions in rising powers of $\hbar$, the so-called WKB development,

$$\psi = e^{\frac{iS}{\hbar}} \left(\psi_0 + \frac{\hbar}{i} \psi_1 + \dots \right) ,$$

investigated the 0-th order approximation $j_0 = (j_0^\mu)$ of the current j associated with the matter field ψ , respectively ψ_0 . Following an idea of J. Audretsch, they proved an interesting property:

Proposition 1 (Audretsch/Gähler/Straumann) *The 0-th approximation of the current j_0 in the WKB-development of a Dirac field or a Klein-Gordon field ψ with mass “function” m (scalar Weyl field of weight -1) on a Weylian manifold with spin structure is geodesic, if and only if the covariant derivative D of m vanishes:*

$$D_{j_0} j_0 = 0 \longleftrightarrow Dm = 0$$

For a **proof**, see the original publication (Audretsch e.a. 1984, 47f.).

Proposing geodesity of the 0-th order of a Dirac or a Klein-Gordon current as a compatibility criterion of Weyl geometry with quantum mechanics, Audretsch and coauthors drew the *striking conclusion that only integrable Weylian manifolds are compatible with quantum physics* (pay regard to their geometrical lemma (3)). Moreover, the natural gauge condition $m \equiv \text{const}$ led them to consider Riemann gauge as the preferred gauge in their context.

In consideration of the result of Audretsch, Gähler and Straumann, only integrable Weyl manifolds seem to be acceptable for our large scale context.¹² We will develop a modified version of what Weyl called an “extension of relativity” inside this restriction (Weyl 1921).

We do not draw the consequence, however, that the gauge condition derived from the Dirac or Klein Gordan equations describes matter behaviour directly. In this respect we keep closer to the view as it is developed in (Canuto e.a. 1977, Tiwari 1989) and continue to respect Weyl’s warning that there may be reasons to assume that matter fields behave as if they adapt to local structure fields (“calibration by adjustment”). These may lead to results different from “calibration by transfer” in the sense of gauge geometry. Although we consider the “short wave limiting” argument of our authors (as they call the 0-th approximation) as compelling with respect to its structural constraints (integrability of the length connection), we do not follow the second step of their evaluation (gauge condition $m \equiv \text{const}$).

If one does not follow this step, it becomes possible to establish a *conservative extension* of GRT, which allows for the possibility that matter properties are represented by a gauge condition different from Riemann gauge, although, of course, the standard assumption of Riemann gauge keeps its validity in most of the standard applications of GRT. The specification “conservative” is here used just like in theory extension in the sense of the foundations of mathematics: In our case, the Einstein equation is lifted into an

¹²In the light of the scale invariance in deep elastic scattering experiments, it may be premature, however, to accept the restriction to integrable Weyl geometry as definitive on all scales. The semi-classical equations used in the proposition apparently lose their meaning in the high energy/small scale domain and demand for “second quantization” or other formal tricks to draw information from them.

extended physico-geometrical framework, but without any change of dynamical properties for the old models. On the other hand, new models can be constructed which were not conceivable before. For the extended theory we sometimes use the abbreviation eGRT.

We start from considering the Einstein equation

$$Ric - \frac{1}{2}\bar{R}g = 8\pi\frac{G}{c^4}T \quad (8)$$

under the aspect of scale gauge in Weyl geometry (Ric the Ricci tensor as above, $\bar{R}$ scalar curvature, T the doubly covariant energy-momentum tensor, G the Newtonian gravitational constant, and c velocity of light).

Remark 6 Let us start with an elementary heuristic observation. The left hand side (l.h.s) of the equation is obviously gauge invariant (cf. remark 5). For the r.h.s. we calculate gauge weights by the elementary assumption that the weights are consistent with the dimensional construction of the quantities. We use simple square brackets $[]$ to denote physical dimensions and double square brackets $[[]]$ for gauge weights of the quantities; l , t , m , E are used as symbols for any length, time, mass, or energy quantity. The Newton constant has dimension $[G] = [l]^3[t]^{-2}[m]^{-1}$. If we want to treat it consistently in a scale gauge geometric theory, it has to be represented by a scalar Weyl field of gauge weight

$$[[G]] = [[l]^3[t]^{-2}[m]^{-1}] = 3 - 2 - (-1) = 2 .$$

The components of the co-contravariant energy-momentum tensor T_j^i is of the same gauge weight as energy density, $[[T_j^i]] = -4$. As we are working with a doubly contravariant version of the Einstein equation, which is exactly the one in which the l.h.s becomes gauge invariant, the right hand side tensor is $T_{ij} = g_{i\mu}T_j^\mu$ and thus of gauge weight $[[T_{ij}]] = -4 + 2 = -2$. Thus the whole r.h.s. is of gauge weight $2 - 2 = 0$. It is composed of scale gauge covariant quantities in such a way, that as a whole it is gauge invariant (as it must be the case, if there shall be a chance for generalization of the equation to gauge geometry).

This shows already by an elementary observation that the Einstein equation ought to be generalizable to Weyl geometry without modification of the underlying dynamical law.

In fact, it is easy to transfer the Hilbert-Einstein variational principle to the case of Weylian manifolds.¹³ We consider a Weylian manifold M with metric $[g, \varphi]$ with integrable length connection φ , W-L-C connection Γ , curvature tensor R , Ricci curvature Ric , and scalar curvature $\bar{R}$. We

¹³This holds even for the non-integrable case although, by the above mentioned reasons, we confine ourselves to the integrable case.

suppose the Langrange density of the gravitational field Γ as given by a gauge invariant density of the form of the Hilbert-Einstein action (equ. (9) below). Because of the gauge weights $[[\bar{R}]] = -2$, $[[\sqrt{|g|}]] = \frac{1}{2}(2 \cdot 4) = 4$, the representing object for the Newton gravitational “constant” has to be a scalar Weyl field of weight $[[G]] = 2$ in order that (9) becomes gauge invariant. We arrive at

Theorem 3 *Let M be an integrable Weyl manifold with metric $[g, \varphi]$, W-L-C connection Γ , scalar curvature $\bar{R}$, and G a nowhere vanishing positive scalar Weyl field of weight 2 on M . Then the Hilbert-Einstein action of the pair (M, G) , defined by*

$$\mathcal{L}_H := \frac{1}{16\pi G} \bar{R} \sqrt{|g|} , \quad (9)$$

is scale-gauge invariant. If, moreover, a scale-gauge invariant Lagrange density of matter and non-gravitational fields $\mathcal{L}_M := L_M \sqrt{|g|}$ is given on M , the variational equation for the combined action

$$\mathcal{S}_H + \mathcal{S}_M := \int \mathcal{L}_H dx + \int \mathcal{L}_M dx$$

is the Einstein equation (8) with an energy momentum tensor given by

$$T_{\mu\nu} := -\frac{1}{2\sqrt{|g|}} \frac{\partial \mathcal{L}_M}{\partial g^{\mu\nu}} . \quad (10)$$

Proof The variational calculation of the Hilbert-Einstein action can be transferred from semi-Riemannian to integrable Weyl manifolds. We norm all the Weylian metrics over which the variation takes place by the condition $G \equiv \text{const}$. That can be done like in the case of the Weyl gauge (lemma (4); here one only has to *divide* by G , $\lambda^2 = G^{-1}$, if G is not yet constant). Now the variational calculation proceeds formally like in the semi-Riemannian case:

$$\begin{aligned} \delta \mathcal{S}_H &= \int \frac{\partial \mathcal{L}_H}{\partial g^{\mu\nu}} \delta g^{\mu\nu} dx \\ \delta \mathcal{S}_M &= \int \frac{\partial \mathcal{L}_M}{\partial g^{\mu\nu}} \delta g^{\mu\nu} dx \end{aligned}$$

Because of the gauge normation for G , we get

$$\begin{aligned} \frac{\partial \mathcal{L}_H}{\partial g^{\mu\nu}} \delta g^{\mu\nu} &= \frac{1}{16\pi G} \frac{\partial(\bar{R} \sqrt{|g|})}{\partial g^{\mu\nu}} \delta g^{\mu\nu} = \frac{1}{16\pi G} \left(\frac{\partial \bar{R}}{\partial g^{\mu\nu}} + \frac{\bar{R}}{\sqrt{|g|}} \frac{\partial \sqrt{|g|}}{\partial g^{\mu\nu}} \right) \sqrt{|g|} \delta g^{\mu\nu} \\ &= \frac{1}{16\pi G} (Ric - \frac{1}{2} \bar{R} g_{\mu\nu}) \sqrt{|g|} \delta g^{\mu\nu} , \end{aligned}$$

just like in the classical case.¹⁴ Note, however, that in this calculation the curvature expressions contain contributions from the length connection φ , if the chosen gauge (called below “gravitational gauge”) is not identical to Riemann gauge. The condition of stationarity,

$$\delta\mathcal{S}_H + \delta\mathcal{S}_M = 0 ,$$

leads directly to the Einstein equation.

The gauge used above will obviously play a preferred role in any attempt of establishing a reasonable semantics for Weyl geometric models in relativity. On the other hand, for each empirical interpretation of a Weyl geometric model of general relativity we have to specify a gauge which coincides with the measurement of atomic clocks and distances measured by material devices. We therefore introduce

Definition 8 *A Weyl geometric model of extended general relativity (eGRT) is given by an integrable Weyl manifold together with an everywhere positive scalar Weyl field G of weight 2. G is called the “gravitational constant”. A gauge for which G is constant in the literal sense ($G(x) \equiv \text{const}$ for all $x \in M$) is called **gravitational gauge** of the model. The gauge which expresses measurements by atomic clocks and material devices will be called **matter gauge**.*

In order to avoid an unnecessary multiplicity of differing gauges, we will assume

$$\text{matter gauge} = \text{gravitational gauge}$$

in the sequel. Of course, this involves the assumption that measurements by material objects lead to one and the same value of the gravitational constant anywhere and at any time in the physical “cosmos”. If we wanted to analyze the consequences of P.A.M. Dirac’s and P. Jordan’s different assumption (of a time dependent gravitational constant) in our framework, we had to give up this identification. That will not be done in this investigation, although the next example shows a weak reflection of Dirac’s idea..

Example 2 Take $G := f^2$ in the semi-Riemannian gauge of a Robertson-Walker-Weyl manifold (example 1). Rescaling by $\lambda = f^{-1}$ gives the gravitational gauge (g_m, φ_m) . Then the Riemannian component of the metric leads to “static rests spaces”,

$$g_m = \left(\frac{d\tau^2}{f} \right) - d\sigma^2, \quad \varphi_m = d \log f.$$

¹⁴Compare, e.g., (Weinberg 1972, 360). Here we use a different sign convention for the energy-momentum tensor.

In our approach (not in Dirac's and Jordan's !) the assumption of a “changing gravitational constant” *in semi-Riemannian gauge* leads to a physically preferred gauge in which G is unchanging. The latter is different to the gauge preferred by purely differential geometrical reasons. Moreover, a reparametrization of the “cosmic time” parameter $t := \int \frac{d\tau}{f}$ simplifies the coordinate expression of the metric,

$$g_m = dt^2 - d\sigma^2, \quad \varphi_m = d \log f.$$

Of course, G is a “true constant” in this gauge (we have chosen it so).

Obviously the Weyl geometric extension of GRT is weak and conservative in the sense that any “classical” general relativistic model of semi-Riemannian geometry remains one in the extension, without any change of its geometrical properties or its dynamics. It is, nonetheless, a veritable *extension* (even though a much more modest one than originally intended by Weyl), because the integration of the Einstein equation allows here a change of scale gauge. Its solution acquires, so to speak, an additional “parameter of integration”.

Because of the restriction to integrable Weyl manifolds, the scale factor and the corresponding scale connection φ are constrained by the differential equation

$$d\varphi = 0. \tag{11}$$

The Einstein equation (8) and the integrability constraint (11) together characterize the physical geometry in our weakly extended general relativity (eGRT).

In this extension, the Einstein equation contains additional terms from the Weylian length connection, if gravitational gauge (matter gauge) (g, φ) is different from Riemann gauge, i.e. $\varphi \neq 0$. The Ricci tensor and scalar curvature of the Weyl-Levi-Civita connection Γ can be decomposed into terms derived from the Riemannian component g only and these additional terms. Let us denote the Ricci tensor and scalar curvature of the Riemannian component g of our gauge by Ric_g and $\bar{R}_g$. Then the l.h.s of the Einstein equation is naturally decomposed into the classical Einstein tensor of the Riemannian component g of the Weylian metric (g, φ) and a term which contains all the contributions from the non-vanishing length connection. We denote the latter by Λ_φ and arrive at a decomposition of the form

$$Ric - \frac{1}{2}\bar{R}g = Ric_g - \frac{1}{2}\bar{R}_g g + \Lambda_\varphi, \tag{12}$$

which can be read as a defining equation for Λ_φ . In our models this additional term plays a role analogous to the cosmological term of standard cosmology. Of course, we can only expect a functional analogy. Λ_φ cannot be brought into the form constant times g . Moreover, it is gauge dependent and can be “gauged away” in Riemann gauge. The functional analogy is strong enough, however, to introduce the following

Terminology 1 *The term Λ_φ of equation (12) will be called the cosmological term of Weyl geometry, or in short the **Weylian cosmological term** of the gauge invariant Einstein equation.*

In fact, Weyl tried at several places to arrive at a more natural derivation of a cosmological term from his gauge geometrical approach. He tried to subsume Einstein’s cosmological term as a “special case” of his and to make Einstein’s term look more “natural” (Weyl 1921, 46). This could not work out by different reasons.

We have already seen that our weak extension of GRT by lifting it to integrable Weyl geometry does *not* imply the introduction of a new dynamical field. We only gain the liberty to investigate models in which Riemann gauge and gravitational gauge (and with it matter gauge) are no longer identical, like in the classical case. So we have new questions to ask for all classical cosmological models with long range extrapolation of geometrical and dynamical behaviour of “the world”: Can we be sure that gravitational gauge and matter gauge are identical with Riemann gauge, even over long distances and into regions of extreme curvature? Or can small modifications of gauge assumptions lead to simpler models in the mathematical or physical sense, although they may coincide with the recieved models in many aspects (hopefully not for all) up to observational error? We shall see in the sequel, that such a tiny modification pays out for cosmological model building.

3. Cosmological redshift and Robertson-Walker-Weyl models

If we want to think cosmology in geometric terms, we have to specify a *cosmic time flow* U and/or an *observer field* X on the manifold M , i.e., an everywhere timelike future oriented vector field on M of unit length. Each such field allows to consider infinitesimal space-time splits, by considering $U^\perp$ (or $X^\perp$) at each $p \in M$ as the infinitesimal rest space at p with respect to the class of (“cosmic”, or more general) observers. Usually one even assumes the existence of a global foliation of M , orthogonal to the field of time flow $U(p)$ at each point $p \in M$, with strong homogeneity and isotropy conditions on the spatial leaves. Moreover, observer and cosmic time flow fields are usually identified, $X = U$. That imposes a natural restriction on the observer field as a *cosmic* or *comoving observer field*.

All this translates to the gauge geometric context without major modifications. We only have to take into account that the unit length condition results in gauging observer fields with weight -1 . We have to consider

$$\begin{aligned} U & \text{ with respect to } (g, \varphi) \text{ s. th. } |U(p)|_g = 1, \\ \tilde{U} & \text{ with respect to } (\tilde{g}, \tilde{\varphi}) \text{ s. th. } |\tilde{U}(p)|_{\tilde{g}} = 1. \end{aligned}$$

Because of $|\tilde{U}(p)|_{\tilde{g}} = \lambda |\tilde{U}(p)|_g$, this implies

$$\tilde{U}(p) = \lambda^{-1} U(p).$$

For a consistent use of gauge geometry in cosmology, a *cosmic time flow field* U or an *observer field* X has thus to be represented by a timelike future oriented Weyl vector field $[X]$ of gauge weight -1 with unit length in every gauge.

Example 3 Take a Robertson-Walker-Weyl manifold $(M, [\tilde{g}, 0])$ as in example 1, with $\tilde{g} = d\tau^2 - (fd\sigma)^2$, and define the observer field in Riemann gauge $(\tilde{g}, 0)$ as given by

$$\tilde{X}(\tau_0) := \frac{\partial}{\partial \tau}|_{\tau_0}.$$

If Riemann gauge is considered as the matter gauge of the model, we arrive at standard cosmology described in the Weyl setting. The choice of any other gauge as matter gauge will lead to new models in Weyl geometry.

Terminology 2 a) A scale integrable Weylian manifold with the specification of a matter gauge (g_m, φ_m) (equivalently gravitational gauge, i.e., $G \equiv \text{const}$) and a time flow (comoving observer) field X of gauge weight -1 (future oriented time-like and unit length) is called a **time flow model** of eGRT. The defining data are $(M, (g_m, \varphi_m), X)$.

b) We will call a model like in example (3) a **Robertson-Walker-Weyl model**. It will be called standard, iff the reference (matter) gauge is the Riemann gauge of the Weylian manifold.

Remark 7 An isomorphism between two time-flow models $(M, (g_m, \varphi_m), X)$ and $(\tilde{M}, (\tilde{g}_m, \tilde{\varphi}_m), \tilde{X})$ is given by an isomorphism of the Weylian manifolds $(\beta, [\lambda])$ (as in remark (2)) and a point-dependent family of Lorentz transformations $\alpha = \alpha(p)$ in the tangent fibres of M , such that

$$\tilde{X}(\beta(p)) = \beta_p^* \alpha(p)(\lambda(p)X(p))$$

(where β^* denotes the operation on tangent vectors etc. induced by the differential of β). Such an isomorphism allows to (Lorentz-) “rotate” the observer field at any point, in other words to transform between different observer fields by “local” Lorentz transformations, besides the application of an isomorphism of Weylian manifolds.

For our investigation, the expression of cosmological redshift by the scale connection of matter gauge will be important. The motion of photons is described by null geodesics, the energy transfer of a photon (respectively the redshift of its frequency) between a point p of emission and a point q of observation is measured with respect to the local space-time split defined by an observer flow X on M (assuming that the emitting oscillator moves with X). The oscillation time T of the photon, its frequency ν and its energy E are related by:

$$E = \hbar\nu, \quad ET = \hbar.$$

Redshift is calculated by

$$z(p, q) = \frac{T'}{T} - 1 = \frac{\nu}{\nu'}.$$

In classical GRT the transfer of photon oscillation time from an emitting system kinematically described by $X(p)$ to an observer described by $X(q)$, respectively of photon energy, is mathematically represented by the orthogonal projection of the tangent of a nullgeodesic on the timelike directions of the comoving observer flow $X(x)$ at p , respectively at q . Parallel transport of tangents along null geodesics is well defined in Weyl geometry by the Weyl-Levi-Civita connection. One might be tempted to calculate the redshift z of a photon passing from p to q in a Weylian manifold along a null geodesic $\gamma(t)$, $0 \leq t \leq 1$, with respect to the observer flow X in eGRT like in GRT. Comparison of quantities derived from the orthogonal projections on a vector field Y on M by $g_{(\gamma(t))}(\gamma'(t), Y(\gamma(t)))$ would not make sense in gauge geometry, however, even for a gauge invariant field Y . One has to take into account the calibration by transfer of the metric g (definition (6)), and compare the recalibrated quantities $l^2(t)g_{(\gamma(t))}(\gamma'(t), Y(\gamma(t)))$. This is a gauge invariant expression (the gauge weights of l^2 and g cancel, as the weight of length transfer in dependence of the endpoint is $[[l]] = -1$, cf. remark (4)).

As the observer (time-flow) field X is of weight -1 , it is natural to give a gauge invariant characterization of redshift in Weyl geometry, which coincides with the classical one in Riemann gauge, in the following way.:

Definition 9 *For a time-flow Weyl model $(M, (g_m, \varphi_m), X)$, let $\alpha(t)$ be defined along any nullgeodesic $\gamma : [0, 1] \rightarrow M$ with $\gamma(0) = p$ and $\gamma(1) = q$ by orthogonal projection on the observer field, corrected by length transfer,*

$$\alpha(t) := l^2(t)g_{\gamma(t)}(\gamma'(t), X(\gamma(t))), \quad (13)$$

where $l(t) := l(\gamma(t))$ is the length transfer function along the geodesic, normed by the condition $l(0) := 1$.

The **redshift function** z of the model is then defined on the subset $Z \subset M \times M$ of nullgeodesically connected pairs of points $(p, q) \in M \times M$ by

$$z(p, q) + 1 := l(1) \frac{\alpha(0)}{\alpha(1)}. \quad (14)$$

With these specifications we get:

Lemma 5 *The redshift function $z(p, q)$ in a time-flow Weyl model is gauge invariant. Its value is identical to the redshift calculated in the semi-Riemannian gauge according to the principles of standard cosmology.*

Proof The gauge invariance is a consequence of cancelling of the gauge weights of the factors:

$$\begin{aligned} [[z + 1]] &= [[l]] - [[\alpha]] = [[l]] - (2[[l]] + [[g]] + [[X]]) \\ &= -[[l]] - [[g]] - [[X]] = +1 - 2 + 1 = 0 \end{aligned}$$

In semi-Riemannian gauge the length transfer is constant, $l \equiv 1$; thus equations (13), (14) reduce to the standard formula in classical relativity.

We have thus arrived at a conservative implementation of redshift in eGRT. Looking at different gauges of physical geometry is the symbolical equivalent of admitting that we do not have direct knowledge about “real” expansion of spatial sections. In fact, such a property is never directly observable and depends on the specific gauge condition imposed (knowingly or unknowingly). In this sense, it is a theoretic construct. Trying to find out different gauges and what they allow to make understandable, corresponds to *adapting our geometrical frame to matter and energy structures the best we can*. It may turn out, that expansion fits best, but it may be that different answers will stand at the end.

Remark 8 In standard relativity, GRT, the wave vector $k(s)$ of a plane wave modelling a photon semiclassically is bound to a null geodesic $\gamma(s)$ by the relation

$$k(s) = C \gamma'(s)$$

(where C is a constant to norm the “emission energy” $E(s_0)$.) Energy decrease (redshift) of a photon has to be expressed by the relation of γ' to the observer field. In classical relativity any kind of energy decrease of photons, including gravitational one (like, e.g., in the Schwarzschild solution), has in this sense a “quasi-kinematical” formal characterization. Although this method offers, for most cases, a well developed and consistent possibility to represent photon energy in differential geometric terms, it is not the only one. In Weyl geometry, it is natural to renorm the wave vector by the length transfer function $l(s)$ along null geodesics,

$$k(s) = C l(s) \gamma'(s). \quad (15)$$

It is even compelling to do so, if one wants to uphold consistency under gauge change. $k(s)$ then becomes a gauge quantity (a Weyl field). As geodesics are defined gauge invariantly, the gauge weight of a wave vector field k along γ is clearly $[[k]] = [[l]] = -1$, which is consistent with the gauge weight of energy. The energy of a photon for an observer with time flow field X develops along γ according to

$$E(p) = g_p(k(s), X(p)) = g_p(l\gamma', X) \quad (16)$$

(setting here $C = 1$). That is consistent with equations (13) and (14),

$$E(p) = lg_p(\gamma', X) = \frac{\alpha(s)}{l(s)} = (z + 1)^{-1}(\gamma(s)).$$

In the expression $E = \alpha(s)l^{-1}(s)$, the first factor $\alpha(t)$ may be considered as a “kinematical contribution” to redshift. It depends essentially on the choice of the observer field and includes space kinematical effects. The second factor, $l^{-1}(t)$, characterizes something like a “gravitational” or, more general, “tired light contribution” to redshift (imputed by the gauge). The quotation marks indicate here that, in general, this distinction is only of formal value. A change of gauge transforms the components into one another, as demanded by the equivalence principle. In a gauge in which the “kinematical contribution” vanishes, $\alpha \equiv 1$, the cosmological redshift appears as completely due to the energy damping expressed by the length transfer $l(t)$. The above equation shows that in this case the photon energy propagates by calibration transfer of weight -1, in the sense of Weyl geometry (definition 7).

This remark may seem abstract and methodological. The investigation of Robertson-Walker-Weyl models shows, however, that we can really calculate with the effects of equivalent exchange between “kinematical” and “gravitational/tired light” contributions to cosmological redshift by an appropriate change of gauge. Used in this way, *Weyl geometry allows to give an enhanced mathematical expression to the equivalence principle.*

Example 4 In a Robertson-Walker-Weyl model

$$M = I \times_f S_\kappa, \quad \tilde{g} : d\tau^2 - (fd\sigma)^2, \quad \tilde{X} = \frac{\partial}{\partial \tau}$$

with *arbitrary* matter gauge (i.e., independent of whether we assume the model to be standard or not), the redshift of a photon emitted at a point $p = (\tau, x)$ and observed at $p_0 = (\tau_0, x_0)$ (with $\tau < \tau_0$ and x, x_0 such that p and p_0 are nullgeodetically connected) is given by

$$z(p, p_0) + 1 = \frac{f(\tau_0)}{f(\tau)}.$$

This follows from standard results on classical Robertson-Walker models and our lemma (5).

For Robertson-Walker models, cosmological redshift depends only on the cosmological “time” parameters of the points of emission and observation. Thus there is a natural extension of the redshift function to all of $M \times M$, given by the r.h.s of the above formula. Keeping the observer p_0 fixed and varying p , we consider $z(p, p_0)$ as a function of p only, $z(p) := z(p, p_0)$. It even

makes sense to write $z(\tau) = z(p_\tau)$ in short, for any p_τ with time coordinate τ . We now look for a gauge in which the length transfer function satisfies

$$l(\tau) = z(\tau) + 1 = \frac{f(\tau_0)}{f(\tau)}.$$

This is achieved by applying the scale gauge factor

$$\lambda(\tau) = \frac{f(\tau_0)}{f(\tau)}$$

to the semi-Riemannian gauge. Then

$$g := \lambda^2 \tilde{g} = f^2(\tau_0) \left(\left(\frac{d\tau}{f} \right)^2 - d\sigma^2 \right), \quad \varphi = -d \ln \lambda = f(\tau_0) d \ln f$$

satisfies the condition. Note that in this gauge, which will be called *Hubble gauge* in a moment, the warping of the fibre metrics is exactly compensated by the rescaling factor $\lambda(\tau) = \frac{f(\tau_0)}{f(\tau)}$. We have thus arrived, in this gauge, at an unwarped static Riemannian component of the Weylian metric like in example 2. In general, the infinitesimal linearization of redshift with timelike distance

$$\varphi(X) = (f' dt) \frac{\partial}{\partial t} = f' \tag{17}$$

is not constant.

E. Hubble was always careful not to claim that he had detected an “expansion” of the observed universe, but only a systematic relationship between redshift and distance. In the gauge above, the redshift is completely encoded in the Weylian length connection φ , independent of the answer to the question, which gauge is “physical”, i.e., the matter gauge. Of course, Hubble left it open to more precise measurements in the future, whether the linearity of the dependence could be extrapolated. In any case, the latter is an interesting theoretical possibility. These are reasons enough to introduce the following

Definition 10 *a) If a time-flow Weyl model $(M, (g_m, \varphi_m), X)$ has a gauge (g, φ) , such that the redshift $z(\gamma(t))$ along any nullgeodesic γ is determined by the length transfer $l(\gamma(t))$ only (i.e., $\alpha = l^2 g(\gamma', X) = \text{const}$),*

$$z(\gamma(t)) + 1 = l(\gamma(t)), \tag{18}$$

*we call (g, φ) the **Hubble gauge** of the model and φ its **Hubble form**.*

b) If for a Hubble gauge of a time-flow model the linearized redshift-time dependence is constant,

$$\varphi(X) = \text{const} =: H, \quad H \in \mathbb{R}^+ \cup \{0\},$$

*the Hubble form will be called **time-homogeneous** and H its **Hubble constant**.*

In the sequel we will use the terminology *cosmological Weyl models* for those time-flow models which possess a Hubble gauge. This is the case for all R-W-W models (last example). Obviously “the” Hubble gauge is determined only up to a (global) constant factor, if it exists. Moreover, it is clear that time-homogeneity results in a constant factor for the linearized redshift-distance relationship *in Hubble gauge only*.

We have now identified three gauges in Robertson-Walker-Weyl models, which are of theoretical interest: Riemann gauge, Weyl gauge, and Hubble gauge. The first two are always different, if warping is not trivial ($f \neq \text{const}$). The last two differ in general, but may coincide in special cases. In the following we establish criteria for their coincidence.

Proposition 2 *In a Robertson-Walker-Weyl cosmology*

$$M = I \times_f S_\kappa, \quad \tilde{g} : d\tau^2 - (fd\sigma)^2, \quad \tilde{\varphi} = 0, \quad \tilde{X} = \frac{\partial}{\partial \tau},$$

with any matter gauge the following holds:

a) *Hubble gauge and Weyl gauge coincide, iff*

$$f''f + f'^2 = \text{const}. \quad (19)$$

b) *In particular Weyl gauge and Hubble gauge are identical, if the Hubble gauge is time homogeneous.*

Proof Let us call λ_W and λ_H the gauge functions from Riemann gauge to Weyl gauge and Hubble gauge respectively (see examples (1) and (4)),

$$\begin{aligned} \lambda_W^2 &= |\tilde{R}| = 6 \left| \left(\frac{f'}{f} \right)^2 + \frac{\kappa}{f^2} + \frac{f''}{f} \right|, \\ \lambda_H &= z(p) + 1 = \frac{\text{const}}{f(\tau)}. \end{aligned}$$

a) We thus find $\lambda_H \sim \lambda_W$, iff

$$\left| \left(\frac{f'}{f} \right)^2 + \frac{\kappa}{f^2} + \frac{f''}{f} \right| \sim \frac{1}{f^2},$$

Multiplication by f^2 leads to (19). The argumentation is invertible.

b) is an immediate result of the definition of time-homogeneity, equation (17), and a).

Remark 9 Hubble and Zwicky suspected that the physical reason for cosmological redshift is not to be looked for in expansion of space. If this is correct, redshift is much better modelled by a Hubble form in the extended

relativistic framework of integrable Weyl geometry than by a formal “expansion factor”, even if reinterpreted by the equivalence principle. In such a case, Hubble gauge has to be accepted as the preferred “physical gauge” of the model, i.e., its matter gauge, rather than the traditional default choice of the Riemann gauge. It is very interesting to notice that the simple assumption of time-homogeneity of Robertson-Walker-Weyl models (which in the cosmological discourse about the middle of the last century used to be called, slightly metaphysically, the “perfect cosmological principle”) leads to the consequence that *Hubble gauge and Weyl gauge are identical*.

If the Zwicky-Hubble assumption is right, we are able to describe the physical geometry of the cosmos by a “Weylianized” Robertson-Walker type structure. If, moreover, the laws of physics (including the reasons for the cosmic tiring of photons) are time-homogeneous, material bodies behave as if they were gauged according to scalar curvature. This need not be considered as a “real” process of physical calibration, but can be understood as a geometrical consequence of the physicality of the Hubble form and of time-homogeneity. The assumption of a “real physical calibration” of material bodies to the scalar curvature was introduced as an interesting, but highly ad-hoc argument by Hermann Weyl during the discussion of the question whether his gauge geometry might be useful for physics, in spite of A. Einstein’s criticism. He came back to this idea in his attempts to apply gauge geometry to cosmological modelling in the fifth edition of his book on *Raum - Zeit - Materie* (Weyl 1923, 299f).

Here we have arrived at the *same formal result* from different basic assumptions.

It thus seems legitimate to call a spatially maximally homogeneous (and therefore also isotropic) Weyl model which possesses a time-homogeneous Hubble form a “Weyl universe”. In the following definition, we take A. Walker’s results into account, which show that, in the semi-Riemannian view, such models can be described by Robertson-Walker manifolds. Moreover we use the simplification of the metric by the Hubble gauge as shown in the last example.

Definition 11 *A cosmological Weyl model $\mathcal{M} := (M, (g, \varphi), X)$ with matter gauge (g, φ) and observer field X in this gauge will be called a **Weyl universe**, iff the following holds:*

(i) *M can be written as a product manifold*

$$M = \mathbb{R} \times S_\kappa,$$

where S_κ is a Riemannian 3-manifold with 3-metric $d\sigma^2$ of constant (ex-ante) sectional curvature $\kappa = \frac{k}{a^2}$ ($k \in 0, \pm 1$, $a \in \mathbb{R}^+$),

(ii) *the Riemannian component of the gauge (g, φ) is given (in geometrical*

units, $c = 1$, and using coordinates $(x_0, x) \in \mathbb{R} \times S_\kappa$) by

$$g : ds^2 = dx_0^2 - d\sigma^2, \quad (20)$$

(iii) the corresponding scale connection is of the simplest non-trivial type

$$\varphi = H dx_0, \quad H \in \mathbb{R}^+ \cup \{0\}, \quad (21)$$

(iv) and the observer field is

$$X := \frac{\partial}{\partial x_0}.$$

In physical units, the Weylian metric contains the velocity of light c , with $c dt = dx_0$, and the Hubble connections acquires the form

$$\varphi = H dx_0 = c H dt = H_0 dt,$$

where $H_0 := cH$ is the usual Hubble constant of astronomers (while our H is often written as $H_1 = c^{-1}H_0$). On S_κ we prefer to use Riemann's coordinates (x_1, x_2, x_3) , because then the metric acquires the well known homogeneous form on spatial sections and shows the close approximation to Euclidean space:

$$d\sigma^2 = \frac{\sum_{\alpha=1}^3 dx_\alpha^2}{(1 + \frac{\kappa}{4} \sum_1^3 x_\alpha^2)^2}.$$

4. Basic properties of Weyl universes

We know from the definition of Hubble gauge and from proposition 1 that the defining gauge (g, φ) of a Weyl universe is its Weyl gauge. Because of integrability, a Weyl universe possesses a Riemann gauge. The scaling function λ from Weyl gauge to Riemann gauge is calculated by integrating along “time-flow” lines $x_0 \in [a, y]$ lifted from the first component $\mathbb{R}$ to M ,

$$\lambda_a(y) = e^{\int_a^y \varphi(\gamma'(\tau)) d\tau} = e^{H(y-a)}.$$

Along curves in spatial sections, λ is constant, therefore the notation $\lambda_a(y)$ is justified. Usually we norm $a := 0$; then we get

$$\lambda(y) := \lambda_0(y) = e^{Hy},$$

or in physical units

$$\lambda(t) = e^{cHt} = e^{H_0 t}. \quad (22)$$

Riemann gauge becomes $\tilde{g} := \lambda^2 g$ and is given by the quadratic form

$$d\tilde{s} = e^{2Hx_0} dx_0^2 - e^{2Hx_0} d\sigma^2.$$

Reparametrization by

$$\tau = H^{-1} e^{Hx_0} \quad (23)$$

(like in example 2) leads to

$$d\tilde{s}^2 = d\tau^2 - (H\tau)^2 d\sigma^2 \quad (24)$$

(of course, here $\tilde{\varphi} = 0$). The observer field is gauged into

$$\tilde{X} = \lambda^{-1} X = e^{-Hx_0} \partial x_0 = \partial \tau$$

(using the abbreviation $\partial x := \frac{\partial}{\partial x}$) and is represented in the new coordinate τ by the same coordinate vector field $(1, 0, 0, 0)$ as X with respect to the x_0 coordinate. We arrive at

Lemma 6 *A Weyl universe with Hubble constant H is a Robertson-Walker-Weyl model with linear warp function*

$$f(\tau) = H\tau,$$

which has Weyl gauge as matter gauge.

In semi-Riemannian gauge, Weyl universes therefore look like the simplest cases of (linearly) “expanding” universes, while in Weyl gauge the Riemannian component of the Weyl metric is a static Lorentz type product of $\mathbb{R}$ by S_κ . According to the different types of sectional curvature we introduce:

Terminology 3 a) For sectional curvatures $\kappa = \frac{k}{a^2}$, $k = \pm 1, 0$ we speak of *Einstein-Weyl universes* in the case $k = +1$, and of *Minkowski-Weyl* or *Lobachevsky-Weyl universe* in the cases $k = 0$ or $k = -1$ respectively.

b) The *module* ζ of a Weyl universe with $H > 0$ is defined by

$$\zeta := \frac{\kappa}{H^2} = \frac{k}{(aH)^2}; \quad (25)$$

a is called the *ex-ante radius of curvature* ($k \neq 0$).

The terminology “module” is chosen because of

Proposition 3 a) *Two Weyl universes with modules ζ and $\tilde{\zeta}$ are isomorphic, iff*

$$\zeta = \tilde{\zeta}.$$

b) *For $H = 0$ there are three isomorphy types, characterized by k .*

Proof a) Consider first the case $k \neq 0$:
“ $\Leftarrow$.” S_{k/a^2} is transformed by a homothety with factor $\eta := \frac{\tilde{a}}{a}$ into $S_{\tilde{k}/\tilde{a}^2}$.
Thus we apply a constant gauge factor $\lambda(x) \equiv \eta$. The length connection remains changed, $\tilde{\varphi} = \varphi - d \ln \lambda = \varphi = H dx_0$. We rescale the observer field by $\lambda^{-1} = \eta^{-1}$,

$$\tilde{X} = \partial \tilde{x}_0 = \eta^{-1} \partial x_0,$$

which implies $d\tilde{x}_0 = \eta dx_0$ and

$$\tilde{\varphi} = H dx_0 = \eta^{-1} H d\tilde{x}_0.$$

We thus arrive at

$$\tilde{H} = \eta^{-1} H = \frac{a}{\tilde{a}} H,$$

and therefore $\zeta = \tilde{\zeta}$.

“ $\Rightarrow$.” If, on the other hand $aH \neq \tilde{a}\tilde{H}$, a homothety necessary to bring the metrics to coincide in spatial directions, does not transform the length connections into each other.

In the case $k = 0$ and any $H > 0$ we rescale by $\lambda = H$. This leads to $\tilde{H} = \tilde{H}^{-1} H = 1$; thus all non-trivial k Minkowski-Weyl universes are isomorphic.

b) For $k = 0, H = 0$, we have the Minkowski space with trivial length connection, $\varphi = 0$. It cannot be transformed isometrically into a form with $H \neq 0$. In this sense, it is degenerate. Similarly for the other cases with $H = 0, k = \pm 1$, the classical Einstein universe and its hyperbolic relative, the static “Lobachevsky universe”.

Remark 10 The non-degenerate Weyl universes ($H \neq 0$) form a continuous 1-parameter space of isomorphy classes, characterized by the module ζ . The Hubble constant H is a (global) scaling quantity. Empirically it has been determined as $H = H_1 = c^{-1} H_0 \approx 2.3 \pm 0.2 (10^4 \text{ Mpc})^{-1}$.¹⁵ For historically minded readers it may be worthwhile to observe that Milne’s “kinematical cosmology” can now be re-read as being “nearly” a Minkowski-Weyl universe with $\zeta = 0$ (a linearly expanding Minkowski space in the Robertson-Walker picture). The whole model class can be seen as a kind of ζ -deformation of this generic object. It is a pity that Milne had no idea of the Weylian metric as a unifying concept;¹⁶ he rather used “two metrics” (corresponding to the Riemannian components of our Weyl respectively Riemann gauges) and preferred the “wrong one” from our point of view. He was fond of the expansionary picture, not only by physical reasons (Kragh 1996, 61, 66).

¹⁵Corresponding to the value $H_0 \approx 70 \pm 7 \text{ kms}^{-1} \text{ Mpc}^{-1} \approx 1.4 \pm 0.1 (10^{10} \text{ Y})^{-1}$.

¹⁶Milne could have had one; all mathematical tools were present at the time. That Weyl universes have not been considered much earlier, seems to be a good case for a “post-mature development” in the sense of J. Stachel (Stachel 2004).

For empirical studies of cosmology in the framework of Weyl universes, it will be crucial to determine an observational value for ζ on the basis of the most reliable data we have and to investigate, whether the result is consistent with the evidence of diverse other classes of observations. It is interesting to see that the supernovae luminosity data of the *Cerro Tololo Inter-American Observatory* group (CTO) and the *Supernova Cosmology Project* (SCP) (Perlmutter e.a. 1999) are as well represented by Weyl universes as in the Friedman-Lemaître model class, although in the latter a rather unnatural second parameter, the cosmological constant, has been introduced into the model space to enhance flexibility.¹⁷

In the present use of F-L models, the cosmological constant is interpreted as (constant) vacuum energy density. Their model space is usually parametrized by $(\Omega_m, \Omega_\Lambda)$, the relative contributions of mass and of the cosmological constant (“vacuum”) to energy density. The curvature contribution Ω_κ is related to the other two by $\Omega_m + \Omega_\Lambda + \Omega_\kappa = 1$. In the Weyl geometric approach, Ω_Λ is a structural effect of the Weylian cosmological term, as we will see in remark 11, and contains no free parameter. Weyl universes do without the symbolical artefact of an adaptable “vacuum energy density” which causes practitioners of cosmology so much headache, because it looks as if it were time-dependent (“dynamical”). Some remarks on such questions will be made in the outlook of this paper; a little more can be found in (Scholz 2004a).

The affine (W-L-C) connection of a Weyl universe can be calculated by Weyl’s theorem (1).¹⁸ If we denote the coefficients of the affine (Levi-Civita) connection on the spatial fibre S_κ by $\tilde{\Gamma}$ and use indexes $\alpha, \beta, \gamma = 1, 2, 3$, we find

$$\Gamma_{\alpha 0}^\alpha = H, \quad \Gamma_{\alpha\alpha}^0 = -g_{\alpha\alpha}H, \quad \Gamma_{\alpha\beta}^\gamma = \tilde{\Gamma}_{\alpha\beta}^\gamma.$$

Denoting the covariant derivative with respect to the Levi-Civita connection in S_κ in direction Y by $\tilde{D}_Y$ we get for the covariant derivatives in M for vector fields Y, Z tangential to the spatial fibres

$$D_{\partial_0}\partial_0 = 0, \quad D_{\partial_0}Y = HY, \quad D_YZ = -g(Y, Z)H + \tilde{D}_YZ.$$

With these expressions we calculate the curvature quantities we are interested in:

Proposition 4 *A Weyl universe, $M = \mathbb{R} \times S$, with Weyl gauge (g, φ) , spatial fibre $S := S_\kappa$ of ex-ante sectional curvature of space sections κ , and $\varphi = Hdx^0$ has Ricci curvature*

$$Ric = 2(\kappa + H^2)d\sigma^2 = -2(\kappa + H^2)g_{\alpha\alpha}(dx^\alpha)^2.$$

¹⁷See the beautiful discussion in (Earman 2001), or also (Demiański 2000).

¹⁸Here, as in other passages where it is adequate, we use the notational conventions of upper and lower indexes for tangent vectors, their linear coefficients, and their duals.

It has an effective sectional curvature in 2-directions tangential to S (i.e., sectional curvature of S embedded in M)

$$\kappa_S = \kappa + H^2$$

and scalar curvature

$$\bar{R} = -6(\kappa + H^2).$$

Components, not stated explicitly, of curvature quantities here given, are 0 (e.g., sectional curvatures containing a timelike direction, etc.).

If doubts should have been left, the expression for the scalar curvature shows again, that the defining gauge of M is Weyl gauge.

As a **corollary**, we calculate the Einstein tensor,¹⁹

$$Ric - \frac{1}{2}\bar{R}g = 3(\kappa + H^2)dt^2 - (\kappa + H^2)d\sigma^2, \quad (26)$$

and the Weylian cosmological term defined in equation (12)

$$\Lambda = 3H^2dt^2 - H^2d\sigma^2. \quad (27)$$

Thus the source term of the Einstein equation has the form of an energy-momentum tensor of an ideal fluid,

$$8\pi \left[\frac{G}{c^4} \right] T = 8\pi \left[\frac{G}{c^4} \right] (\rho dt^2 + p d\sigma^2) \quad (28)$$

(square bracket terms for values in physical units, in geometrical units $c = G = 1$), although with negative (repulsive) pressure. The general form is no surprise, as in any Robertson Walker manifold the r.h.s. of the Einstein equation acquires the form of the energy-momentum of a “strange” fluid ((O’Neill 1983)). The important, and from the standard view surprising point is that constant values for energy density and pressure can go in hand with redshift. That justifies the announcement of this result as:

Theorem 4 *The source term of the Einstein equation of a Weyl universe with ex-ante sectional curvature κ and Hubble constant H is an energy-momentum tensor of a strange²⁰ ideal fluid (equ. (28)). In gravitational gauge (Weyl gauge), energy density ρ and pressure p are constant and have the values*

$$\rho = \frac{3}{8\pi} \left[\frac{c^4}{G} \right] (\kappa + H^2), \quad p = -\frac{1}{8\pi} \left[\frac{c^4}{G} \right] (\kappa + H^2).$$

¹⁹The note (Scholz 2004a) contains a sign error in the $d\sigma^2$ - term

²⁰“Strange” fluid, because of negative pressure.

Remark 11 The crucial difference between the statical models of cosmology and Weyl universes is expressed by the Hubble form and the Weylian cosmological term (27). If these are more than formal gadgets, they represent the geometrical aspects of a tired light/gravitational modification of the transfer of wave-vectors along null geodesics (compare remark 8). In such a case, one can reasonably expect that the matter-field Lagrangian $\mathcal{L}_M$ which gives rise to the r.h.s. of the Einstein equation (10) splits into a matter term $\mathcal{L}_m$ proper and a term expressing the tired light “mechanism” (which may very well be quantum physical) $\mathcal{L}_\Lambda$,

$$\mathcal{L}_M = \mathcal{L}_m + \mathcal{L}_\Lambda.$$

In analogy to how standard cosmology interprets its cosmological term as “vacuum energy”, we can then split up the r.h.s. of the Einstein equation of Weyl universes into a “mass term” ρ_m and a “cosmological term” ρ_Λ of the energy density:

$$8\pi \left[\frac{G}{c^4} \right] \rho_m = 3\kappa \quad \text{and} \quad 8\pi \left[\frac{G}{c^4} \right] \rho_\Lambda = 3H^2.$$

It is important to keep in mind that the Weylian cosmological term *cannot be considered as “vacuum contribution”* in the sense of the standard approach, as it is not of the form *const* · *g*. Its contribution to the r.h.s. of the Einstein equation has the form of a strange fluid, like the matter term proper. It could therefore, in principle, just as well be assimilated to the latter. That also allows to consider the possibility that only the energy density of the Weylian cosmological term is due to a physical structure lying at the base of the Hubble redshift (if different from the matter content of the universe), while the pressure term makes sense only in the overall balance ($p = p_m + p_\Lambda$).

If we use the standard definition of critical energy density, which also for Weyl universes corresponds to $\kappa = 0$,

$$\rho_{crit} = \frac{3}{8\pi} \left[\frac{c^4}{G} \right] H^2,$$

the relative contributions of the mass term $\Omega_m := \frac{\rho_m}{\rho_{crit}}$ and of the cosmological term $\Omega_\Lambda := \frac{\rho_\Lambda}{\rho_{crit}}$ become

$$\Omega_m = \frac{\kappa}{H^2} = \zeta \quad \text{and} \quad \Omega_\Lambda = \frac{H^2}{H^2} = 1.$$

These relations are just another expression of the beautiful correspondences between sectional curvature values in spatial directions and energy densities, which have already been established. We rephrase them here in geometrical units:

$$8\pi\rho = 3\kappa_S, \quad 8\pi\rho_m = 3\kappa, \quad 8\pi\rho_\Lambda = 3H^2.$$

That is, the total energy density determines the ex-post sectional curvature in spatial directions of a Weyl universe, matter density determines the ex-ante curvature, and the energy density of the Weylian cosmological term determines the curvature contribution of the Hubble connection.

Remark 12 The source term of Weyl universes satisfies the state equation

$$p = -\frac{1}{3}\rho. \quad (29)$$

At first glance one may have the impression that the equilibrium problem raised by Eddington for the Einstein universe, or even worse Einstein’s gravitational collapse problem, might reappear in a slightly modified form. This is not the case, however, as Weyl universes do not rely on the classical cosmological term. Moreover, we now know that star and galaxy matter (luminous matter ρ_{lum} with proportionality factor $\Omega_{lum} = \frac{\rho_{lum}}{\rho_{crit}} \approx 0.005$) makes up only a very small part of the matter content of the universe. The main bulk of material content of the universe seems to be ionized gas, plasma (most of it ionized hydrogen H_1), mixed with some neutral gas and dust components. Internal repulsion in the intergalactic medium, arising from electromagnetic forces and, perhaps, elementary particle physics components, have to be taken into account. Their overall action may very well balance Einstein’s gravitational attraction which, taken for itself, would let “the world” appear to be damned to collapse.

In our context, there is no reason for bringing exotic matter into play. “Exotic matter” has been postulated as an ad-hoc hypothesis to avoid refutation of the theory of primordial nucleosynthesis by empirical data on dynamical mass density and to alleviate the problems arising in standard cosmological theories of structure formation. The dynamically determined mass density ρ_d , with relative value compared to the critical density $\Omega_d \approx 0.3$, is now at least one order of magnitude higher than the maximal density of baryonic matter, allowed by the hypothesis of primordial nucleosynthesis, $0.04 \leq \Omega_{bar} \leq 0.05$. No indications for exotic new types of matter have been found in several decades of intense research. It is time to accept that the *primordial nucleosynthesis hypothesis has been refuted* in its core (baryon density) by empirical evidence, although certain features of it (high energy contribution to nucleosynthesis) will probably survive in a modified way.²¹

Thus we simply assume that dynamical matter is dark (non-luminous) ordinary matter, dominated by ionized gas, comparable to the one in intergalactic space, only more concentrated around galaxies, clusters, and superclusters. Although the dynamics of the highly diluted ionized matter (apparently a *plasma* in the sense of H. Alfvén, A. Peratt e.a.) has still to be understood, it is clear that the old game of contradicting attractive and

²¹For a short hint, see outlook.

repulsive forces comes back into the game, now as the interplay of gravitational and electrodynamical forces.²² Cosmic space has to be understood as constituted in the interplay of such attractive and repulsive forces of matter. If Weyl universes turn out to be empirically satisfying models of cosmological observations on a broader range, we even can *conclude* that there is dynamical equilibrium in “the cosmic mean” (i.e., mean values formed over very large regions of space-time), with equation (29) as a global expression for such an equilibrium. The overall interplay of attraction and repulsion in the compound system of intergalactic plasma (including cosmic ray contributions), dust, luminous matter, etc., has to satisfy a state equation of a “strange” fluid. In allusion to an idea of Greek natural philosophers, I propose to call it the *hyle* equation. This term does *not impute* new kinds of matter.

5. Redshift, light cone and luminosity decrease

The defining gauge of Weyl universes is Hubble gauge, it is therefore immediate to write down the redshift function. For a photon with trajectory γ , emitted at $p = (t, x)$ and observed at $q = (t_0, x_0)$ ($t < t_0$), redshift is given, according to equ. (18), by

$$z(p, q) + 1 = e^{\int_t^{t_0} \varphi(\gamma'(\tau)) d\tau} = e^{H_0(t_0 - t)}.$$

That agrees, of course, with the value expected from the Robertson-Walker picture (c.f. equs. (23), (24)),

$$z + 1 = \frac{H\tau_0}{H\tau} = e^{H_0(t_0 - t)}.$$

Comparison with equ. (22) shows

$$z + 1 = \lambda^{-1}.$$

Here, the metrical perspective is completely different to the Robertson-Walker picture. From the perspective of Weyl gauge, the blow up of redshift, close to the “initial singularity” of the Riemann gauge, appears as the result of a metrically deformed view of infinity. The same holds for other quantities dependent on the cosmic time parameter. As Weyl gauge is the physically dominant perspective (matter gauge) in our new models, with literally constant G and constant energy densities, *this shift of perspective must not be dismissed as only a formal effect* of reparametrization,

In particular, cosmic time distances, measured in matter gauge (more precisely, in the Riemannian component of the Weyl metric of matter gauge) go to infinity with redshift:

$$\Delta t := t_0 - t = H_0^{-1} \ln(z + 1).$$

²²For first interesting steps, see (Peratt 1991, Peratt 1995a, Verschuur 1995).

Because it is such an important property, we formulate the standardized form of the redshift development with time/distance as

Proposition 5 *In a Weyl universe, the redshift-time relation for a photon emitted at $t < 0$ and observed at $t_0 = 0$ (“today”) is given by*

$$z(t) = e^{H_0|t|} - 1 = \lambda^{-1} \leftrightarrow |t| = H_0^{-1} \ln(z + 1), \quad (30)$$

where λ is the scaling function from Weyl to Riemann gauge, normed by the condition $\lambda(0) = 1$.

From standard results on Robertson-Walker manifolds (O’Neill 1983, 355f.) and by changing coordinates as well as gauge, one can determine explicit expressions for null geodesics in Weyl gauge. This allows to “verify” (i.e. here, to check the consistency of definitions and methods) the relations between null geodesics, wave vector, length connection, and redshift given above in definition 9 and the equations of example 8. Here we content ourselves with more general observations on the light cone and its geometry.

Because of conformal invariance of the light cone, the *traces* of null geodesics in Weyl universes with Weyl gauge (g, φ) are identical to traces of null geodesics of the Riemannian component g only. They can easily be described. Let us denote by r the length of the projection of the segment $t \leq s \leq t_0$ of a null geodesic $\gamma(s)$ into the spatial standard fibre S_κ . Abbreviating the redshift between t and t_0 by z , we find

$$r = c|t - t_0| = H^{-1} \ln(z + 1).$$

We call the intersection of the light cone with spatial fibres a *light sphere* and r its *radius*. The area of light spheres and volumes of light cone segments can easily be calculated.²³

Lemma 7 *In a Weyl universe of ex-ante sectional curvature κ , Hubble constant H , and module $\zeta = \kappa H^{-2}$, the following holds:*

a) *The area $O = O_\kappa$ of light spheres in dependence of redshift is given by*

$$O_\kappa = \frac{4\pi}{\kappa} SIN_k^2(\sqrt{\zeta} \ln(z + 1)), \quad \left(O_0 = \frac{4\pi}{H^2} \ln^2(z + 1) \right),$$

where $k = \pm 1$ or 0 and $SIN_1 := \sin$, $Sin_0 := id$, $SIN_{-1} := \sinh$.

b) *The volume $V(z_1, z_2)$ scanned by the light cone between redshift values $z_1 \leq z_2$ is*

$$V(z_1, z_2) = \int_{z_1}^{z_2} (z + 1)^{-1} O(z) dz. \quad (31)$$

²³Warning: In (Scholz 2004a, 5) the area formula for Riemann gauge are given erroneously in place of those in Weyl gauge. The magnitude formula on the same page is correct.

The **proof** of the first statement is obvious, as we calculate areas of spheres with radius $r = H^{-1} \ln(z + 1)$ in the 3-geometries of constant curvature radius $a = \kappa^{-\frac{1}{2}}$ and use (25). For b), we only need to integrate over infinitesimal volume layers Odx^0 of the light cone, substitute $dx^0 = cd|t|$, and use equation (30).

The decrease of luminosity of (electromagnetic) radiation sources with redshift results from a damping of the energy flow by two reasons, the change with increasing z of the area of light spheres (“normally”, but not always, an increase), and a damping of energy quantities due to length transfer of energy flux, or more precisely of calibration by transfer, in Weyl geometry. In the standard expanding models, the latter effect is interpreted as a reduction of flux etc. because of expansion of spatial sections. Here we use a natural principle for the application of scale gauge to energy quantities:

Principle 1 *Propagation of energy quantities of gauge weight k in the “cosmic ether”, i.e., the compound system of the gravitational and the electromagnetic fields, is mathematically represented by calibration transfer of gauge weight k with respect to Hubble gauge.*

Example 5 a) Photon energy E , respectively the wave vector k is a scalar Weyl field of gauge weight $[[E]] = -1$ and propagates along null geodesics by calibration transfer of weight -1 in *Hubble gauge* (see remark 8). This is no surprise; we have defined Hubble gauge with this property in mind.
b) The energy flux F of radiation of astronomical objects is observed as energy per time and area. Thus F has the gauge weight $[[F]] = [[E]][[t]]^{-1}[[l]]^{-2} = -1 - 1 - 2 = -4$. According to the principle, it propagates by calibration transfer of weight -4 .

This is a natural assumption in Weyl geometry. One may compare it to the view of standard cosmology, where a damping factor $(1 + z)^{-2}$ is assumed for the energy flux, one factor $(1 + z)^{-1}$ for redshift, another one due to time dilation for comoving observers. Surface brightness and intensity are calculated with a damping factor $(1 + z)^{-4}$ (Peterson 1997, 152f.).

On this basis we can calculate the magnitude-redshift dependence in Weyl universes.

Theorem 5 *Assuming principle 1 for the propagation and measurement of electromagnetic energy in a Weyl universe,*

a) *the energy flux $F(z)$ of a radiating cosmic source depends on redshift z and the area of light spheres $O(z)$ measured in Weyl gauge like*

$$F(z) \sim (z + 1)^{-4} O(z)^{-1}.$$

b) After redshift correction of the measured energy by the factor $(z + 1)$, the corrected energy flux F_{corr} is proportional to

$$F_{corr} \sim (z + 1)^{-3} O(z)^{-1}. \quad (32)$$

Proof a) We consider transmission of radiation from the source at a point $p = (t, x)$ to a point at $p_0 = (t_0, x_0)$ along a null geodesic γ . According to principle 1 energy flux F propagates by calibration transfer of gauge weight -4 . This damping of the energy is only due to the effect of the transfer according to the Hubble form, respectively the gauge factor λ from Weyl to Riemann gauge as its integrated version. Here we consider calibration depending on the origin of the path rather than on its endpoint; thus we have to take the reciprocal (see equ. 7), and the flux is proportional to λ^4 ,

$$F(\gamma(t)) \sim \lambda^4(\gamma(t)).$$

From equ. (30) we know $\lambda = (z + 1)^{-1}$; therefore the contribution of the calibration transfer to damping is

$$F(z) \sim (z + 1)^{-4}.$$

In addition, radiation energy emitted at (t, x) during a short time interval (an “infinitely short” one) is distributed over the whole light sphere at the spatial section lying over t_0 , with area $O(z(t, t_0)) =: O(z)$. That results in an area contribution to damping according to

$$F(z) \sim O(z)^{-1}.$$

If we neglect absorption by cosmic matter, we arrive at an overall effect in Weyl geometry

$$F(z) \sim (z + 1)^{-4} O(z)^{-1}.$$

The correction in b) is direct.

Remark 13 Readers who are doubtful of principle 1, may want to avoid calibration by transfer and prefer to calculate in the Robertson Walker picture with traditional methods of scaling down the flux in “expanding universes” (not to forget the gauge factors between Weyl and Riemann gauge). The result should be the same. In any case, the empirical data of supernovae Ia magnitudes are a beautiful empirical test for the physical acceptability of the principle *in the frame* our approach (see below, outlook).

In the sequel we refer to the redshift corrected values, F_{corr} , if not explicitly stated otherwise, because the redshift-energy correction as in part b) of the theorem is usually applied by astronomers during the K -correction for apparent magnitudes (Goenner 1994, 80).

The apparent magnitude of a cosmic source is defined by

$$m := -2.5 \log F + C, \quad (33)$$

where the constant C contains the dependence on the *absolute magnitude* M , i.e., the magnitude in which the same source would appear to a terrestrial observer, if it was placed at the norm distance of classical astronomy $d_0 := 10\text{pc} = 10^{-5}\text{Mpc}$ (just in reach of good parallax measurements). Moreover, C depends on the cosmological assumptions in each model, in particular on the value for the Hubble constant H_0 . With the values for F from the last theorem we get

$$m = -2.5 \log(z+1)^{-3} O(z)^{-1} + C = 5 \log(z+1)^{\frac{3}{2}} \sqrt{O(z)} + C,$$

and with lemma 7:

Corollary *In a Weyl universe with module ζ and spatial ex-ante curvature type $k = -1, 0, +1$, the redshift-corrected apparent luminosity m_k of a cosmic radiation source is:*

$$m_k(z) = 5 \log \left(\frac{(z+1)^{\frac{3}{2}}}{\sqrt{|\zeta|}} \text{SIN}_k(\sqrt{|\zeta|} \ln(z+1)) \right) + C, \quad (34)$$

($\text{SIN}_1 = \sin$, $\text{SIN}_0 = \text{id}$, $\text{SIN}_{-1} = \sinh$, as above). For $k = 0$, this is to be understood in the sense of the limit $\zeta \rightarrow 0$,

$$m_0 = 5 \log \left((z+1)^{\frac{3}{2}} \ln(z+1) \right) + C.$$

Sometimes it is useful to substitute the constant C in the formulae of the corollary by an expression in the absolute magnitude at redshift $z_0 = e^{Hd_0} - 1 \approx H d_0$,

$$C = M - 5 \log \left(\frac{e^{1.5 H d_0}}{\sqrt{\zeta}} \text{SIN}(\sqrt{\zeta} H d_0) \right) \approx M - 5 \log(H d_0).$$

Then we get an explicit function for $m_k(z, M, \zeta, H)$, depending on the redshift, the absolute magnitude, the module ζ of the Weyl universe (or Ω_m), and the Hubble constant. The last approximation is so good that it can be taken as a “numerical equality” for all practical purposes.

If we accept the source term interpretation of the Einstein equation as in remark 11, $\zeta = \Omega_m$, we get a “more physical” version of (34). For $k = 1$, it acquires the form

$$m_k(z) = 5 \log \left(\frac{(z+1)^{\frac{3}{2}}}{\sqrt{\Omega_m}} \sin(\sqrt{\Omega_m} \ln(z+1)) \right) + C. \quad (35)$$

The great simplicity of Weyl universes results in simple explicit expressions for their geometrico-physical properties. As a last example, the angular size of objects with diameter b in distance d can easily be calculated from spherical trigonometry for $k = 1$, because the light cone structure is conformally invariant under deformation of the classical Einstein universe to the Einstein-Weyl one. Angular sizes of such objects are given by

$$\sin \frac{\alpha}{2} = \frac{\sin \frac{b}{2a}}{\sin \frac{d}{a}}, \quad (36)$$

where a is the ex-ante radius of curvature of spatial sections, $a = \kappa^{-\frac{1}{2}} = \frac{H^{-1}}{\sqrt{\zeta}}$.

6. Outlook

Before we come to the end, we want at least to indicate how well the new models behave with respect to observational data. We start with the recent *luminosity-redshift measurements of supernovae Ia* by the *Cerro Tololo Inter-American Observatory* group (CTO) and the *Supernova Cosmology Project* (SCP) (Perlmutter e.a. 1999). These data, complemented by those of the group around A. Riess, have been interpreted as striking evidence in favour of the Friedman-Lemaître model with non-vanishing cosmological constant (Grøn 2002). Together with constraints derived from the “inflationary” hypothesis they lead to the presently accepted standard density parameters $(\Omega_m, \Omega_\Lambda) \approx (0.3, 0.7)$. It is therefore quite remarkable that in our framework *the supernovae data can be fitted just as well by Weyl universes* (figure 1). With $H \approx 2.3 \cdot 10^{-4} \text{ Mpc}^{-1}$ ($H_0 \approx 70 \text{ kms}^{-1} \text{ Mpc}^{-1}$) and $M \approx -19.3$ for the absolute magnitude of supernovae Ia, the best fit for the module of Weyl universes is

$$\zeta \approx 1, \quad \text{with residual dispersion } \sigma \approx 0.318 \approx 1.24\sigma_{data}.^{24}$$

This result can be considered as a first empirical answer to the question, whether the hypothesis of luminosity decrease according to calibration by transfer in Weyl geometry (principle 1) makes sense or not. *It clearly does.*

The fit quality remains very stable under change of ζ . If we relax only a little bit to $\sigma \leq 1.3\sigma_{data}$, we get a wide interval for admissible curvature parameters, including all three curvature types $k = \pm 1, 0$

$$\zeta \in [-0.1, 3.6] =: I_{SNIa}.$$

For $\zeta = 0.3$, the residual dispersion is still only $\sigma_{0.3} \approx 0.325$. Because of the possible identification $\Omega_m = \zeta$ (remark 11), this value is of particular interest, if we want to adapt the model to present data of dynamical mass

²⁴The fit quality for the standard model is $\sigma_{stand} \approx 0.309$ for the parameter choice $(\Omega_m, \Omega_\Lambda) = (0.3, 0.7) \approx 1.21\sigma_{data}$ (my calculation, E.S.). The difference is not significant.




Figure 1: Magnitude-redshift relation in Einstein-Weyl universe, $\zeta = 1$ (no graphical distinction to $\zeta = 1.5$), $H_0 = 70$, $M = -19.3$, compared with recent SN_{Ia} data of the SCP and CTO groups (dots)

density. From the empirical point of view, the supernovae luminosities and the mass density criterion clearly single out the *Einstein-Weyl universes*, $k = +1$, among the whole model class.

Other important data for testing cosmological models are *quasar frequencies* in dependence of redshift. The best values presently available are those of the *Sloan Digital Sky Survey* (first data release) (Fan e.a. 2003). The colleagues of the SDSS have measured 16 713 objects up to $z \leq 5.4$ and gave a concentrated survey in a histogram with 69 intervals of width $\Delta z = 0.076$. The frequencies show a clear and well-formed increase up to about $z \approx 2$ and a rapid decrease for $z > 2.1$. It is very interesting to see that the peak of observed quasar frequencies about $z \approx 2$ corresponds closely to the maximum in volume scanned by the light cone (in equal z -intervals) in Einstein-Weyl universes (equ. (31)).²⁵

A calculation of the numbers of objects to be expected under the assumption of equal distribution in Einstein-Weyl universes for $\zeta = 0.3, 0.8, 1, 1.5$ and 2 gives a surprisingly good agreement of the geometry of the light cone with the observed quasar frequencies up to the peak at $z \approx 2$. For larger values of z , observed quasar frequencies break down too abruptly to be accounted for by geometrical reasons only (figure 2).²⁶ In fact the rapid decline of observed quasar numbers is strongly enforced by the reduction of sensibility of the CCD detectors for $z > 2$ and, perhaps, other selection effects.

Already a rough comparison of the data shows that a parameter interval

$$\zeta \in I_{QF} := [0.3, 2]$$

²⁵Because of the inflationary dogma ($k = 0$) and the expanding geometry, this effect is suppressed in the standard approach.

²⁶Densities of objects have been calculated from respective volumes in Einstein-Weyl universes and SDSS data for $z \leq 2.24$, shortly after the peak.




Figure 2: Frequencies of equally distributed objects in Einstein-Weyl universe with $\zeta = 0.3$ (coarsest dashing), $\zeta = 0.6$ (coarse dashing), $\zeta = 1$ (medium dashing), $\zeta = 1.5$ (undashed), and $\zeta = 2$ (finest dashing), compared with quasar frequencies of SDSS (dots); interval width $\Delta z = 0.076$

fits in with quasar frequencies up to $z \approx 2$. For larger values of z , it leads to an acceptable agreement only for $\zeta \geq 0.6$ and if sufficiently strong selection effects are at work. Which range of the interval I_{QF} is to be preferred, depends on an estimation of selection effects and a comparison of the reliability of other data which are taken into account.

The determination of *dynamical mass density* close to galaxies, clusters, and superclusters leads to an actual estimation of $0.1 \leq \Omega_m \leq 0.5$ for smaller structures and of $0.2 \leq \Omega_m \leq 1$ for larger ones, with a preferred value close to $\Omega_m \approx 0.3$ (Carroll 2001). That is consistent with supernovae and quasar data, but probably not the best we can get. We may reasonably expect a rising value for Ω_m with increasing awareness for dynamical mass in larger regions.

Mass density estimations are precarious data. The best measured cosmological data are those of the *cosmic microwave background* (CMB) with its precise Planck profile and a radiation temperature $T_{CMB} \approx 2.726$ °K with error less than 0.1% (after subtraction of the dipole moment of motion against the CMB rest system).

In Einstein-Weyl universes, a beautiful alternative to the standard explanation of the CMB (as the redshifted view upon a “surface of last scattering”) is possible by a theorem of I.E. Segal. Segal has proved that the quantized Maxwell field on the Einstein universe has a (maximal entropy) equilib-

rium state with radiation of exact Planck characteristic (Segal 1983). The proof translates to the Weyl case because of conformal invariance of the field equations. It is now the question, whether (or how) the Segal background is related to the cosmic background radiation.

In an Einstein-Weyl universe, the Segal background plays the role of a vacuum ground state for the electromagnetic field. We may thus reasonably *conjecture*:

(*) *The energy density ρ_Λ of the Weylian cosmological term is due to the energy density ϵ_{Segal} of the Segal background, $\rho_\Lambda = \epsilon_{Segal}$.*

If this is true, ϵ_{Segal} is equal to the critical density (remark 11), $\rho_{crit} \approx 5.2 \text{ keV cm}^{-3}$, and corresponds to a Planck temperature $T_{Segal} \approx 32^\circ \text{K}$. Although that may appear unlikely at first glance, a second thought shows that any direct identification of such a background equilibrium state will be difficult or even impossible. Everything, including each part of the measuring device, will be immersed in a diffuse bath of the 32°K radiation, which suppresses its discernibility as incoming radiation. Moreover, its spectral range lies deep inside the interval of the cosmic infrared background which is very difficult to measure anyhow (Dwek/Hauser 2001). Thus there seems to be no chance for a *direct* empirical identification of the emission component due to the Segal background.




Figure 3: Refocussing of luminosity (here for a source of absolute magnitude $M = -16$) in Einstein-Weyl universe with $\zeta = 1.5$ (undashed) at the first conjugate point, $z \approx 12$, in comparison with luminosity of an equally strong source in the standard cosmological model ($\Omega_m, \Omega_\Lambda = (0.3, 0.7)$) (dashed)

On the other hand, the radiation characteristics of the region close to the conjugate point of a general observer in an Einstein-Weyl universe would be strongly dominated by the (redshifted) Segal background. Therefore the signal from this region would have a clearly discernible Planck characteristic, be nearly ideally isotropic, and would be highly refocussed at the point of

observation (see figure 3). A sharp delimitation of refocussing about the redshift at the conjugate point suppresses a smearing out of the Planck characteristic of the source, which would otherwise happen. The signal will be corrupted only by interaction along the passage through the spatial sphere S_κ with hot gas around clusters and superclusters, or a slightly increased energy density of the Segal background in these regions. We have to look at empirical indicators which allow to decide whether the conjecture (*) and its consequences are realistic or not.

If the CMB is due to emission of the Segal background in the vicinity of “our” conjugate point, an easy calculation shows that the redshift at the conjugate point has to be

$$z_{conj} = \frac{T_{Segal}}{T_{CMB}} \approx \frac{32}{2.73} \approx 12.$$

That corresponds to a parameter value $\zeta \approx 1.5 \pm 5\%$ of the Einstein-Weyl universe and *stands in pleasing agreement with quasar frequencies and SNIa luminosities*. We may consider this agreement as a first positive empirical test for the conjecture. Of course more of them are needed, most important a detailed analysis of anisotropy data from this point of view, before we can claim more than a hypothetical plausibility for it.

On the theoretical level, it is worthwhile to notice that, in the frame of Weyl universes and under the assumption of the correctness of (*), *the two empirically most important cosmological parameters, temperature T_{CMB} of the microwave background and Hubble constant H , determine ζ and with it the physical mean geometry of the cosmos in the large without ambiguity, in principle, and with high precision in practice.*²⁷ If other observations support the conjecture, we will attribute these data much more credibility than mass density estimations from dynamical observations, which cannot account for the diluted mass in intercluster regions anyhow.

If the Segal background gives a realistic explanation of the cosmic microwave background (with or without conjecture (*)), the anisotropies arise naturally as a result of the passage through material inhomogeneities, in particular cosmic plasma (“hot gas”) concentrated around clusters, cluster groups, and superclusters, or through regions of slightly increased temperature of the Segal background. The most important contributions to anisotropies with multipole moment $l \approx 200$ are expected to be caused by objects on the supercluster level in the segment of the spatial S_κ with $1 \leq z \leq 6$, which covers 80 % of the total volume until the first conjugate point. For $\zeta = 1.5$, objects of diameter 40 *Mpc* in this redshift range appear under an angular size between 0.7 deg and 1 deg (figure 4). and lead to a peak of the anisotropy signal at a multipole moment of about $l \approx \frac{180}{0.8} = 225$.

²⁷The small error bound $\pm 5\%$ for the estimation of ζ arises from the fourth root to be drawn from the energy density and thus from H^2 .




Figure 4: Angular size of objects with diameter 40 Mpc in Weyl universe with $\zeta = 1.5$

It is very encouraging that astronomers have started to investigate the possibility of an influence of clusters and higher order structures on CMB anisotropy. Recent evaluations of WMAP anisotropy data by a group around T. Shanks (Myers/Shanks e.a. 2004) have demonstrated that there exists, in fact, a *strong anti-correlation between temperature and cluster positions for redshifts up to $z \approx 0.2$* . The authors explain their observation by inverse Compton scattering of photons in hot gas in clusters and cluster groups (“Sunyaev-Zeldovich effect”). They consider their results as a corruption of a more original anisotropy signal stipulated in the standard picture to result from some legendary “acoustic waves” in the primordial soup at the end of the “inflationary phase”. Shanks e.a. expect that with a continuation of the evaluation to $z \approx 0.5$ the “contamination” of anisotropy data, as they call it, may rise considerably.

From our point of view, that *has* to happen. We even can quantify the degree of “contamination”, which is to be expected. It is closely related to the volume increase covered by the light cone in respective z -intervals. The interval $0 \leq z \leq 0.2$ covers 0.23% of the total volume (up to the first conjugate point) in a Weyl universe of $\zeta = 1.5$. Contributions to the anisotropy signal with $l \approx 200$ from redshifts $0.1 \leq z \leq 0.2$ are caused by objects on the cluster level (up to 10 Mpc). The main anisotropy signal is expected to be caused by superclusters with $1 \leq z \leq 6$, as indicated above. Taking the number of partial systems (clusters in superclusters) into account, the anisotropy contribution of this redshift range should be an order of magnitude larger than its volume ratio. Thus it should account for about 1 – 3 % of the whole anisotropy signal of the CMB. For the next interval, $0.2 \leq z \leq 0.5$, envisaged by the group around T. Shanks, the volume proportion rises to 2.5%. Anisotropies with $l \approx 200$ result here from objects in cluster group size. We therefore expect a contribution to CMB anisotropies of about

10 %.

It would be very welcome if the estimation of the anisotropy contributions could be continued until $z \leq 1.5$ or even $z \leq 2$. For $z > 0.8$, structures on the supercluster level come into the play. For these redshifts, our volume estimations let us expect a contribution to the whole anisotropy signal of 35 % and 43 %, respectively. A detailed investigation of the CMB anisotropies may even be able to shed light on the origin of the “tired light mechanism”, if this is in fact the reason for the good modelling behaviour of Weyl universes. The reduction of photon energy during the passage of clusters (or higher order structures) may have the same general reason as the Hubble redshift itself. A photon-photon interaction with the Segal background, slightly increased in energy/temperature in regions of luminous matter concentrations may be an additional cause (“Freundlich-Vigier-Segal effect”), or even an alternative, to the Sunyaev-Zeldovich effect. In this case we had traced a common physical basis for both, the Hubble redshift and the tiny anisotropies impressed on the microwave background signal (or at least parts of them).

Other points have to be discussed with the necessary care and in greater detail than can be done here. Most important among them are those related to the observation that expansion may only be due to a geometrical fiction of the model class which has been accepted as *the* symbolical a-priori during the last decades (F-L models). If physics of the cosmos is better characterized by Weyl gauge, expansion appears as a merely formal feature of Riemann gauge. In fact, in the standard approach the dependence of Ω_Λ on the cosmic time parameter and the famous “accelerated expansion” of the universe suggest that all this may just be a model artefact, without direct (realistic) physical meaning. The behaviour of the “dynamical dark energy” in the standard approach hints even more strongly in this direction.

The attempts of achieving a satisfying model of structure formation by an interplay of repulsive dynamical dark energy plus cosmic expansion and gravitational attraction have run into an impasse. Even under the auxiliary assumption of fictitious gravitational material (exotic dark matter), it was only possible to imitate either large scale structures (“hot dark matter”) or small scale structures (“cold dark matter”), but never both together (Ostriker/Steinhardt 2003). From these heroic attempts we can draw the lesson that in the standard picture the repulsive force is given a much too schematic expression to allow for sufficient flexibility, or even (structural) “creativity” of matter. We therefore have reasons to look forward for progress in the plasma physical alternative program which tries to understand structure formation by an interplay of gravitational attraction with electromagnetic attraction and repulsion inside the ionized cosmic medium. That will surely be easier without the expansionary paradigm than within it.²⁸

²⁸Cf. footnote 22. It will be a challenging task for the history and philosophy of science to have a closer look at the cultural conditions which let it appear plausible to two

Metallicity values of galaxies and quasars give another occasion for a critical review of the idea of a global evolution correlated with expansion. Even though very detailed and long range data are now available, they do not show a systematic correlation between redshift and metallicity (Pagel 1997). Apparently, this lack is not seen as a problem in the actual discussion. To the surprise of astronomers and astrophysicists, however, the metallicity of the quasars with the highest redshifts observed at present shows no indication of lower overall values than quasars and galaxies closer by (Corbin e.a. 2003). Adding to the surprise, very metal poor galaxies of small redshift have been detected (Brinkmann e.a. 2003). *The metallicity of galaxies and quasars turns out to depend much more on local conditions* (most importantly the mass of the object) *and regional evolution histories than on a “global evolution”* clear signs of which are even missing. In the present discourse, the observational facts are explained by diverse exceptional clauses to a general rule which seems to exist in the theoretical expectation of the standard approach only.²⁹

Finally, the empirical indicators of physical states of the cosmos “close to the initial singularity” are no longer as convincing as they appeared to be some decades ago. As mentioned above, the hypothesis of *primordial nucleosynthesis* has been factually refuted by the observational data on dynamical mass and the lack of success for diverse attempts to find a substitute in strange (“exotic”) matter (remark 12). Recent high precision lattice gauge calculations of the assumed electro-weak phase transition (which has been considered as the clue to baryogenesis in the “early universe” in the standard approach) complement this picture. The calculations demonstrate the existence of a *theoretical upper bound* of 74 GeV for a Higgs-boson mass compatible with electro-weak phase transition (Csikor e.a. 1999). This bound is inconsistent with the present *experimental lower limit* of 89.8 GeV for the Higgs mass. Because of reliable error estimates for the theoretical and for the experimental result, *the combination of the standard model of elementary particle physics (SM) with the standard approach to cosmology (SC) has lead to another anomaly for the hypothesis of a “primordial phase of the world”*.³⁰

With primordial nucleosynthesis and the electroweak phase transition, two important features of a hypothetical early phase of the expanding world picture have shown to be wrong. A distributed locus for high energy contributions to nucleosynthesis, which has to exist in addition to stellar nucle-

generations of scientists that a paradigm of nuclear explosion, or of a (post-) “inflationary boost”, can serve as a clue to structure *formation* (rather than destruction) — contrary to the modelling experience of several decades.

²⁹I thank E. Grebel for information on this point.

³⁰The authors of the study comment their result in clear words: “This also means that the SM baryogenesis in the early Universe is ruled out” (Csikor e.a. 1999, 24). By SM the authors refer to the standard model of elementary particle physics. They consider their result as an argument for the necessity for a physics beyond SM.

osynthesis, has entered observability in the form of active galactic nuclei and quasars. Empirical and theoretical knowledge on these structures is growing rapidly. We may expect that they provide more realistic candidates for high energy nucleosynthesis (beyond stellar processes) than the hypothetical primordial phase of the universe in the old paradigm, although new features like a cracking of higher nuclei into light ones enter here and change the overall function of the process (“recycling” of baryonic matter rather than its “genesis”). The Weyl geometric approach provides an alternative without the necessity to invent repair measures inside the paradigm of a “hot initial phase of the universe”.

If, in one way or another, the most important signatures of a global evolution of *the cosmos as a whole* turn out to be unreliable, we have to content ourselves for scientific investigations with observing local, or better, regional evolution of *substructures in the cosmos* and to understand them the best we can. The grand narrations of *The Primordial Phase of The World* can be handed over again to the myths of all the different cultures on our earth. This is no real loss, by the way; questions relating to the evolution of the universe as a whole have always been a part of speculative natural philosophy and continue to be so. In the extremes, they even may be considered as either “meaningless” or as an attempt to approach “the mind of God” etc., depending on taste and the metaphysical perspective we assume. Somewhere in between stand the most recent quantum cosmological speculations of late modernity.

Here, we could only scratch at the surface of these difficult and multifarious problems. For the moment, we have to leave it with first glimpses into a vast and challenging terrain. If Weyl universes appear useful and theoretically attractive to other researchers, there will be better occasions for more detailed discussions of the open questions and, hopefully, some new and unexpected answers.

Acknowledgements: I have to thank many people from different fields for criticism and support in different stages of the research for this project. A talk of P. Cartier in September 2001 was important to me to take up these investigations. In the early stage, H. Goenner and J.H. Eschenburg contributed critically to conceptual questions; H. Arndt and P. Feuerstein gave computational support. During the course of the development of the approach I profited from remarks by K.-H. Kampert, Z. Fodor, D. Lind, W. Rhode, A. Rendall, T.Scholz, E. Grebel, D. Puetzfeld and an anonymous referee of an earlier version submitted to (and rejected by) *Annalen der Physik*. Matthias Kreck was the first mathematician, who took the enterprise seriously; he invested time and enthusiasm in discussions on the mathematics and supported me to find ways to improve and to communicate it to others. Among physicists, the discussions at Freiburg with D. Giulini and H. Roemer gave important hints to me and the interest of J. Renn, J. Stachel and G. Ellis provided a kind form of encouragement. Outside the professional world, it was a particular pleasure for me that Sven Heinze, Sonja Heinze, and Angelika Scheuffele responded

with sympathy and curiosity in the questions dealt with here.

References

- Assis, A.K.T.; Neves, M.C.D.. 1995. "The redshift revisited." *Astrophysics and Space Science* 227: 13–24. Reprinted in (Peratt 1995*b*, 13–24).
- Audretsch, Jürgen; Gähler, F. ; Straumann Norbert. 1984. "Wave fields in Weyl spaces and conditions for the existence of a preferred pseudo-Riemannian structure." *Communications in Mathematical Physics* 95:41–51.
- Brinkman, Jonathan; Fukugita, Masataka; Grebel, Eva; Kniazev, Alexei; Hao, Lei. 2003. "New metal-poor galaxies: The youngest in the universe?" *Astrophysical Journal Letters* August 20. [Abstract <http://www.sdss.org/new/releases/20030818.metapoor.html>. Visited November 10 2003].
- Canuto, V; Adams, P.J.; Hsieh, S.-H.; Tsiang, E.. 1977. "Scale covariant theory of gravitation and astrophysical application." *Physical Review D* 16:1643–1663.
- Carroll, Sean. 2001. "The cosmological constant." *Living Reviews in Relativity* 4: 1–77. [Also in arXiv:astro-ph/0004075v2. Visited November 10, 2003.]
- Corbin, Michael; Freudling, Wolfram; Korista, Kirk. 2003. Iron emission in $z \approx 6$ QSOs and its possible implications. In *Neutron Stars in Supernova Remnants*. ASP Conference Series, vol. 9999. [arXiv:astro-ph/0311043, visited April 25, 2004.]
- Csikor, F.; Fodor, Zoltán; Heitger, J.. 1999. "End point of the hot electroweak phase transition." *Physical Review Letters* 82:21–24.
- Demiański, M. 2000. "History of the cosmological constant." *Annalen der Physik* 9:3–5, 278–287.
- Deser, S. 1970. "Scale invariance and gravitational coupling." *Annals of Physics* 59:248ff.
- Dirac, Paul A.M.. 1973. "Long range forces and broken symmetries." *Proceedings Royal Society London A* 333:403–418.
- Dwek, Eli; Hauser, Michael. 2001. "The cosmic infrared background: Measurements and implications" 5:1–11. [arXiv:astro-ph/0105539 v1 30 May 2001. Visited December 26, 2003.]
- Earman, John. 2001. "Lambda: the constant that refuses to die." *Archive for History of Exact Sciences* 55:189–2001.
- Ehlers, Jürgen; Pirani, Felix; Schild, Alfred. 1972. The geometry of free fall and light propagation. In *General Relativity, Papers in Honour of J.L. Synge*, ed. Lochlainn O’Raifeartaigh. Oxford: Clarendon Press pp. 63–84.
- Ellis, George F.R.. 1989. The expanding universe: A history of cosmology from 1917 to 1960. In *Einstein and the History of General Relativity*, ed. Howard, Don; Stachel, John. Basel/Boston: Birkhäuser, 367–431.
- Ellis, George F.R.; Maartens, R.; Nels, S.D.. 1979. "The expansion of the universe." *Monthly Notices Royal Astronomical Society* 184:439–465.

- Fan, Xiahui; Strauss, Michael e.a. 2003. "The Sloan Digital Sky Survey Quasar Catalog II. First data release". [arXiv:astro-ph/0308443 v1, Aug 2003. Visited November 10, 2003.]
- Folland, George B. 1970. "Weyl manifolds." *Journal of Differential Geometry* 4:145–153.
- Goenner, Hubert. 1994. *Einführung in die Kosmologie*. Heidelberg etc.: Spektrum Verlag.
- Goenner, Hubert. 2004. "On the history of unified field theories." *Living Reviews in Relativity* . [http://relativity.livingreviews.org/Articles/lrr-2004-2, visited June 2, 2004].
- Grøn, Ø. 2002. "A new standard model of the universe." *European Journal of Physics* 23:135–144.
- Hehl, Friedrich; Mc Crea, J. Dermott; Mielke, Eckehard. 1989. Weyl spacetimes, the dilation current, and creation of gravitating mass by symmetry breaking. In *Exact Sciences and Their Philosophical Foundations*., ed. Arnold Oberschelp Volker Weidemann Wolfgang Deppert, Kurt Hübner. Frankfurt/Main: Peter Lang pp. 241–310.
- Hentschel, Klaus.1994. "Erwin Finlay Freundlich and testing Einstein's theory of relativity." *Archive for History of Exact Sciences* 47:143–201.
- Hentschel, Klaus. 1998. *Zum Zusammenspiel von Instrument, Experiment und Theorie*. Rotverschiebung im Sonnenspektrum und verwandte spektrale Verschiebungseffekte von 1880 bis 1960. Hamburg.: Verlag Dr. Kovač.
- Kragh, Helge. 1996. *Cosmology and Controversy: The Historical Development of Two Theories of the Universe*. Princeton: University Press.
- Manin, Yuri. 1981. *Mathematics and Physics*. Basel: Birkhäuser.
- Milne, E.A.. 1935. *Relativity, Gravitation and World-Structure*. Oxford: Clarendon.
- Myers, Adam D.; Shanks, Thomas; Outram, Phillip J.; Frith, William J.; Wolfendale, Aarnold W.. 2004. "Evidence for an extended SZ effect in WMAP data". To appear in *Monthly Notices Royal Astronomical Society*. [arXiv:astro-ph/0306180 v2, visited August 27, 2004]
- O'Neill, Barrett. 1983. *Semi-Riemannian Geometry with Applications to Relativity*. New York: Academic Press.
- Ostriker, Jeremiah, Steinhardt, Paul. 2003. "New light on dark matter." *Science* 300:1909–1913.
- Pagel, Bernhard. 1997. *Nucleosynthesis and Chemical Evolution of Galaxies*. Cambridge: University Press.
- Pecker, J.-C.; Robert, A.-P.; Vigier, Jean-Pierre. 1972. "Sur une interprétation possible du déplacement vers le rouge des raies spectrales dans le spectre des objets astronomiques." *Comptes Rendus Académie des Sciences* 274 B:765–768.
- Peratt, Anthony L. (ed.). 1991. *Physics of the Plasma Universe*. Berlin etc.: Springer.

- Peratt, Anthony L.. 1995*a*. “Plasma and the universe: Large scale dynamics, filamentation, and radiation.” *Astrophysics and Space Science* 227:97–108. Reprinted in (Peratt 1995*b*, 97–108).
- Peratt, Anthony L. (ed.). 1995*b*. *Plasma Astrophysics and Cosmology*. The Second IEEE International Workshop, Princeton, New Jersey, May 10–12, 1993. Dordrecht etc.: Kluwer. Reprint of *Astrophysics and Space Science*, vol. 27, nos. 1–2.
- Perlmutter, S. ; Aldering G; Goldhaber G. e.a. 1999. “Measurement of Ω and Λ from 42 high-redshift supernovae.” *Astrophysical Journal* 517:565–586.
- Peterson, Bradley. 1997. *An Introduction to Active Galactic Nuclei*. Cambridge: University Press. [Introductory chapter also under <http://nedwww.ipac.caltech.edu/level5/Cambridge/frames.html>. Visited November 10, 2003.]
- Puetzfeld, Dirk. 2004. “Status of non-Riemannian cosmology.” Preprint Köln. [<http://arXiv.org/gr-qc/0404119>, visited, June 29, 2004].
- Scholz, Erhard. 2004*a*. “An outline of Weyl geometric models in cosmology.” Preprint Wuppertal . [<http://arXiv.org/astro-ph/0403446>, visited June 2, 2004].
- Scholz, Erhard. 2004*b*. “Herman Weyl’s analysis of the “problem of space” and the origin of gauge structures.” *Science in Context* 17:165–197.
- Segal, Irving E. 1983. “Radiation in the Einstein universe and the cosmic background.” *Physical Review D* 28:2393–2401.
- Stachel, John. 2004. “[Postmaturity].” Preprint Boston.
- Tiwari, Suresh. 1989. “Action principle in Weyl geometry.” *Physics Letters A* 135:315–319.
- Tiwari, Suresh. 2003. “Electron in the Einstein-Weyl space.” [arXiv:gr-qc/03037079, visited August 28, 2004].
- Varadarajan, V.S. 2003. Vector bundles and connections in physics and mathematics: some historical remarks. In *A Tribute to C. S. Seshadri (Chennai, 2002)*, ed. V. Balaji; V. Lakshmibai. Trends in Mathematics Basel etc.: Birkhäuser pp. 502–541.
- Verschuur, Gerrit L.. 1995. “Interstellar neutral hydrogen filaments at high galactic latitudes and the Bennet pinch.” *Astrophysics and Space Science* 227:187–198. Reprinted in (Peratt 1995*b*, 187–198).
- Weinberg, Stephen. 1972. *Gravitation and Cosmology : Principles and Applications of the General Theory of Relativity*. New York etc.: Wiley.
- Weyl, Hermann. 1918. “Reine Infinitesimalgeometrie.” *Mathematische Zeitschrift* 2:384–411, GA II, 1–28, [30].
- Weyl, Hermann. 1921. “Über die physikalischen Grundlagen der erweiterten Relativitätstheorie.” *Physikalische Zeitschrift* 22:473–480. In GA II, 229–236 [46].
- Weyl, Hermann. 1923. *Raum - Zeit - Materie*, 5. Auflage. Berlin: Springer.

- Weyl, Hermann. 1929. "Elektron und Gravitation." *Zeitschrift für Physik* 56:330–352. In (Weyl 1968, III, 245–267) [85].
- Weyl, Hermann. 1930. "Redshift and relativistic cosmology." *London, Edinburgh and Dublin Philosophical Magazine* 9:936–943. GA III, 268–279 [89].
- Weyl, Hermann. 1968. *Gesammelte Abhandlungen, 4 vols.* Ed. K. Chandrasekharan. Berlin etc.: Springer.
- Zwicky, F. 1929. "On the possibilities of a gravitational drag of light." *Physical Review* 33:1623f..